

Revisiting Algebra and Complexity of Inference in Graphical Models

Siamak Ravanbakhsh, University of Alberta
Russell Greiner, University of Alberta

This paper studies the form and complexity of inference in graphical models using the abstraction offered by algebraic structures. In particular, we broadly formalize inference problems in graphical models by viewing them as a sequence of operations based on commutative semigroups. We then study the computational complexity of inference by organizing various problems into an *inference hierarchy*. When the underlying structure of an inference problem is a commutative semiring – i.e., a combination of two commutative semigroups with the distributive law – a message passing procedure called belief propagation can leverage this distributive law to perform polynomial-time inference for certain problems. After establishing the NP -hardness of inference in any commutative semiring, we investigate the relation between algebraic properties in this setting and further show that polynomial-time inference using distributive law does not (trivially) extend to inference problems that are expressed using more than two commutative semigroups. We then extend the algebraic treatment of message passing procedures to survey propagation, providing a novel perspective using a combination of two commutative semirings. This formulation generalizes the application of survey propagation to new settings.

1. INTRODUCTION

Many complicated systems can be modeled as a graphical structure with interacting local functions. Many fields have (almost independently) discovered this: graphical models have been used in bioinformatics (protein folding, pedagogy trees, regulatory networks), neuroscience (formation of associative memory and neuroplasticity), communication theory (low density parity check codes), statistical physics (physics of dense matter and spin-glass theory), image processing (inpainting, stereo/texture reconstruction, denoising and super-resolution), compressed sensing, robotics (particle filters), sensor networks, social networks, natural language processing, speech recognition, combinatorial optimization and artificial intelligence (artificial neural networks, Bayesian networks). Two general perspectives have emerged from these varied approaches to local computation, namely variational [Wainwright and Jordan 2007] versus algebraic [Aji and McEliece 2000] perspectives on graphical models. These two perspectives are to some extent “residuals” from the different origins of research in AI and statistical physics.

In the statistical study of physical systems, Boltzmann distribution relates the probability of each state of a physical system to its energy, which is often decomposed due to local interactions [Mézard et al. 1987; Mézard and Montanari 2009]. These studies often model systems at the thermodynamic limit of infinite variables and the average behaviour through the study of random ensembles. Inference techniques with this origin (e.g., mean-field and cavity methods) are asymptotically exact under these assumptions. Most importantly these studies have reduced inference to optimization through the notion of free energy –a.k.a. the variational approach.

In contrast, graphical models in the AI community have emerged from the study of knowledge representation and reasoning under uncertainty [Pearl 1988]. These advances are characterized by their attention to the theory of computation and logic [Bacchus 1991], where interest in computational (as opposed to analytical) solutions has motivated the study of approximability, computational complexity [Cooper 1990; Roth 1993] and the invention of inference techniques such as belief propagation that are efficient and exact on tree structures. But, most relevant to the topic of this paper, these studies have lead to algebraic abstractions in modeling systems that allow local computation [Shenoy and Shafer 1990; Lauritzen and Jensen 1997].

The common foundation underlying these two approaches is information theory, where the derivation of probabilistic principles from logical axioms [Jaynes 2003] leads to notions such as entropy and divergences that are closely linked to their physical counter-parts *i.e.*, entropy and free energies in physical systems. At a less abstract level, [Yedidia and Freeman 2001] showed that inference techniques in AI and communication are attempting to minimize (approximations to) free energy (also see [Aji and McEliece 2001; Heskes 2003]).

Another exchange of ideas between the two fields was in the study of critical phenomenon in random constraint satisfaction problems by both computer scientists and physicists [Fu and Anderson 1986; Mitchell et al. 1992; Monasson et al. 1999]. Satisfiability is at the heart of the theory of computation and is used to investigate reasoning in AI. On the other hand, the study of critical phenomena and phase transitions is central in the statistical physics of disordered systems. This culminated when a variational analysis lead to the discovery of survey propagation [Mezard et al. 2002] for constraint satisfaction, which significantly advanced the state-of-the-art in solving random satisfiability problems.

Despite this convergence, variational and algebraic perspectives are to some extent complementary – *e.g.*, the variational approach does not extend beyond (log) probabilities, while the algebraic approach cannot justify application of message passing to graphs with loops. This paper is concerned with the algebraic approach. We organize and generalize the previous work on the algebra of graphical models and give several new results on the complexity and limit of inference in this framework. To this end, section 2 broadly formalizes (and extends) the problem of inference using factor-graphs and commutative semigroups. Section 3 organizes a subset of inference problems into an inference hierarchy with increasing levels of computational complexity under PSPACE. Section 4 reviews the distributive property that make efficient inference possible (using Belief Propagation), establishes the difficulty of inference in general commutative semirings and derives a negative result regarding the application of the distributive law beyond commutative semirings. Section 5 moves beyond Belief Propagation and introduces an algebraic interpretation of survey propagation that generalizes its application to new settings.

2. THE PROBLEM OF INFERENCE

We use commutative semigroups to both define what a graphical model represents and also to define inference over this graphical model. The idea of using structures such as semigroups, monoids and semirings in expressing inference has a long history [Lauritzen and Jensen 1997; Schiex et al. 1995; Bistarelli et al. 1999]. Our approach, based on factor-graphs [Kschischang and Frey 2001] and commutative semigroups, generalizes a variety of previous frameworks, including Markov networks [Clifford 1990], Bayesian networks [Pearl 1985], Forney graphs [Forney Jr 2001], hybrid models [Dechter and Larkin 2001], influence diagrams [Howard and Matheson 2005] and valuation networks [Shenoy 1992].

In particular, the combination of factor-graphs and semigroups that we consider here generalizes the plausibility, feasibility and utility framework of [Pralet et al. 2007], which is explicitly reduced to the graphical models mentioned above and many more. The main difference in our approach is in keeping the framework free of semantics (*e.g.*, decision and chance variables, utilities, constraints), that are often associated with variables, factors and operations, without changing the expressive power. These notions can later be associated with individual inference problems to help with interpretation.

Definition 2.1. A **commutative semigroup** is a pair $\mathcal{G} = (\mathcal{Y}^*, \otimes)$, where $\mathcal{Y}^*$ is a set and $\otimes : \mathcal{Y}^* \times \mathcal{Y}^* \rightarrow \mathcal{Y}^*$ is a binary operation that is (I) associative: $a \otimes (b \otimes c) = (a \otimes b) \otimes c$ and (II) commutative: $a \otimes b = b \otimes a$ for all $a, b, c \in \mathcal{Y}^*$. A **commutative monoid** is a commutative semigroup plus an identity element 1 such that $a \otimes 1 = a$. If every element $a \in \mathcal{Y}^*$ has an inverse a^{-1} (often written $\frac{1}{a}$), such that $a \otimes a^{-1} = 1$, the commutative monoid is an **Abelian group**.

Here, the associativity and commutativity properties of a commutative semigroup make the operations invariant to the order of elements. In general, these properties are not “vital” and one may define inference starting from a *magma*.¹

Example 2.2. Some examples of semigroups are:

- The set of strings with the concatenation operation forms a semigroup with the empty string as the identity element. However this semigroup is not commutative.
- The set of natural numbers $\mathcal{N}$ with summation defines a commutative semigroup.
- Integers modulo n with addition defines an Abelian group.
- The power-set 2^S of any set S , with intersection operation defines a commutative semigroup with S as its identity element.
- The set of natural numbers with greatest common divisor defines a commutative monoid with 0 as its identity. In fact any semilattice is a commutative semigroup [Davey and Priestley 2002].
- Given two commutative semigroups on two sets $\mathcal{Y}^*$ and $\mathcal{Z}^*$, their Cartesian product is also a commutative semigroup.

Let $\underline{x} = (x_1, \dots, x_N)$ be a tuple of N discrete variables $x_i \in \mathcal{X}_i$, where $\mathcal{X}_i$ is the finite domain of x_i and $\underline{x} \in \mathcal{X} = \mathcal{X}_1 \times \dots \times \mathcal{X}_N$. Let $I \subseteq \mathcal{N} = \{1, 2, \dots, N\}$ denote a subset of variable indices and $\underline{x}_I = \{x_i \mid i \in I\} \in \mathcal{X}_I$ be the tuple of variables in $\underline{x}$ indexed by the subset I . A factor $f_I : \mathcal{X}_I \rightarrow \mathcal{Y}_I$ is a function over a subset of variables and $\mathcal{Y}_I = \{f_I(\underline{x}_I) \mid \underline{x}_I \in \mathcal{X}_I\}$ is the range of this factor.

Definition 2.3. A **factor-graph** is a pair $(\mathcal{F}, \mathcal{G})$ such that

- $\mathcal{F} = \{f_I\}$ is a collection of factors with collective range $\mathcal{Y} = \bigcup_I \mathcal{Y}_I$.
- $|\mathcal{F}| = \text{Poly}(N)$.
- f_I has a polynomial representation in N and it is possible to evaluate $f_I(\underline{x}_I) \forall I, \underline{x}_I$ in polynomial time.
- $\mathcal{G} = (\mathcal{Y}^*, \otimes)$ is a commutative semigroup, where $\mathcal{Y}^*$ is the closure of $\mathcal{Y}$ w.r.t. $\otimes$.

The factor-graph compactly represents the expanded (joint) form

$$q(\underline{x}) = \bigotimes_I f_I(\underline{x}_I) \quad (1)$$

Note that the connection between the set of factors $\mathcal{F}$ and the commutative semigroup is through the “range” of factors. The conditions of this definition are necessary and sufficient to 1) compactly represent a factor-graph and 2) evaluate the expanded form, $q(\underline{x})$, in polynomial time. In the following we make a stronger assumption that ensures a factor has a tractable tabular form – that is $|\mathcal{X}_I| = \text{Poly}(N)$, which means

¹A magma [Pinter 2012] generalizes a semigroup, as it does not require associativity property nor an identity element. Inference in graphical models can be also extended to use magma (in definition 2.3). For this, the elements of $\mathcal{Y}^*$ and/or $\mathcal{X}$ should be ordered and/or parenthesized so as to avoid ambiguity in the order of pairwise operations over the set. Here, to avoid unnecessary complications, we confine our treatment to commutative semigroups.

$f_I(\underline{x}_I)$ can be explicitly expressed for each $\underline{x}_I \in \mathcal{X}_I$ as an element of $|I|$ -dimensional array or table.²

$\mathcal{F}$ can be conveniently represented as a bipartite graph that includes two sets of nodes: variable nodes x_i , and factor nodes I . A variable node i (note that we will often identify a variable x_i with its index “ i ”) is connected to a factor node I if and only if $i \in I$ – i.e., I is a set that is also an index. We will use ∂ to denote the neighbours of a variable or factor node in the factor graph – that is $\partial I = \{i \mid i \in I\}$ (which is the set I) and $\partial i = \{I \mid i \in I\}$.

Also, we use Δi to denote the **Markov blanket** of node x_i – i.e., $\Delta i = \{j \in \partial I \mid I \in \partial i, j \neq i\}$.

Example 2.4. figure 1 shows a factor-graph with 12 variables and 12 factors. Here $\underline{x} = (x_i, x_j, x_k, x_e, x_m, x_o, x_r, x_s, x_t, x_u, x_v, x_w)$, $I = \partial I = \{i, j, k\}$, $\underline{x}_K = \underline{x}_{\{k, w, v\}}$ and $\partial j = \{I, V, W\}$. Assuming $\mathcal{G}_e = (\mathbb{R}, \min)$, the expanded form represents

$$q(\underline{x}) = \min\{f_I(\underline{x}_I), f_J(\underline{x}_J), \dots, f_Z(\underline{x}_Z)\}.$$

Now, assume that all variables are binary – i.e., $\mathcal{X} = \{0, 1\}^{12}$ and $q(\underline{x})$ is 12-dimensional hypercube, with one value per each assignment at each corner. Also assume each of the factors count the number of non-zero variables – e.g., for $\underline{z}_W = (1, 0, 1) \in \mathcal{X}_W$ we have $f_W(\underline{z}_W) = 2$. Then, for the complete assignment $\underline{z} = (0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1) \in \mathcal{X}$, it is easy to check that the expanded form is $q(\underline{z}) = \min\{2, 0, 1, \dots, 1\} = 0$.

A marginalization operation shrinks the expanded form $q(\underline{x})$ using another commutative semigroup with binary operation $\oplus$. Inference is a combination of an expansion and one or more marginalization operations, which can be computationally intractable due to the exponential size of the expanded form.

Definition 2.5. Given a function $q : \mathcal{X}_J \rightarrow \mathcal{Y}$, and a commutative semigroup $\mathcal{G} = (\mathcal{Y}^*, \oplus)$, where $\mathcal{Y}^*$ is the closure of $\mathcal{Y}$ w.r.t. $\oplus$, the marginal of q for $I \subset J$ is

$$q(\underline{x}_{J \setminus I}) \stackrel{\text{def}}{=} \bigoplus_{\underline{x}_I} q(\underline{x}_J) \quad (2)$$

where $\bigoplus_{\underline{x}_I} q(\underline{x}_J)$ is short for $\bigoplus_{\underline{x}_I \in \mathcal{X}_I} q(\underline{x}_{J \setminus I}, \underline{x}_I)$, which means to compute $q(\underline{x}_{J \setminus I})$ for each $\underline{x}_{J \setminus I}$, one should perform the operation $\oplus$ over the set of all the assignments to the tuple $\underline{x}_I \in \mathcal{X}_I$.

We can think of $q(\underline{x}_J)$ as a $|J|$ -dimensional tensor and marginalization as performing $\oplus$ operation over the axes in the set I . The result is another $|J \setminus I|$ -dimensional tensor (or function) that we call the **marginal**. Here if the marginalization is over all the dimensions in J , we denote the marginal by $q(\emptyset)$ instead of $q(\underline{x}_\emptyset)$ and call it the **integral** of q .

Now we define an inference problem as a sequence of marginalizations over the expanded form of a factor-graph.

Definition 2.6. An **inference problem** seeks

$$q(\underline{x}_{J_0}) = \bigoplus_{\underline{x}_{J_M}}^M \bigoplus_{\underline{x}_{J_{M-1}}}^{M-1} \dots \bigoplus_{\underline{x}_{J_1}}^1 \bigotimes_I f_I(\underline{x}_I) \quad (3)$$

²Important factor-graphs that violate this assumption are the ones with high-order sparse factors [Tarlow and Zemel 2010; Potetz and Lee 2008]. Although it is possible to obtain polynomial time message passing updates for special high-order factors, general high-order factors do not have polynomial representation.




Fig. 1. A factor-graph with variables as circles and factors as squares. The partitioning of variables indicated by the shading in parts defines an inference problem over this factor-graph.

where

- $\mathcal{Y}^*$ is the closure of $\mathcal{Y}$ (the collective range of factors), w.r.t. $\bigoplus_1, \dots, \bigoplus_M$ and $\otimes$.
- $\mathcal{G}_m = (\mathcal{Y}^*, \bigoplus_m)$ $\forall 1 \leq m \leq M$ and $\mathcal{G}_e = (\mathcal{Y}^*, \otimes)$ are all commutative semigroups.
- $J_0, \dots, J_L$ partition the set of variable indices $\mathcal{N} = \{1, \dots, N\}$.
- $q(\underline{x}_{J_0})$ has a polynomial representation in N — i.e., $|\mathcal{X}_{J_0}| = \text{Poly}(N)$

Note that $\bigoplus_1, \dots, \bigoplus_M$ refer to potentially different operations as each belongs to a different semigroup. When $J_0 = \emptyset$, we call the inference problem **integration** (denoting the inquiry by $q(\emptyset)$) and otherwise we call it **marginalization**. Here, having a constant sized J_0 is not always enough to ensure that $q(\underline{x}_{J_0})$ has a polynomial representation in N . This is because the size of $q(\underline{x}_{J_0})$ for any individual $\underline{x}_{J_0} \in \mathcal{X}_{J_0}$ may grow exponentially with N (e.g., see claim 3.3). In the following we call $\mathcal{G}_e = (\mathcal{Y}^*, \otimes)$ the expansion semigroup and $\mathcal{G}_m = (\mathcal{Y}^*, \bigoplus_m)$ for each m in $\{1, \dots, M\}$ is a marginalization semigroup.

Example 2.7. Going back to example 2.7, the shaded region in figure 1 shows a partitioning of the variables that we use to define the following inference problem:

$$q(\underline{x}_{J_0}) = \max_{\underline{x}_{J_3}} \sum_{\underline{x}_{J_2}} \min_{\underline{x}_{J_1}} \min_I f_I(\underline{x}_I)$$

We can associate this problem with the following semantics: we may think of each factor as an agent, where $f_I(\underline{x}_I)$ is the payoff for agent I , which only depends on a subset of variables $\underline{x}_I$. We have adversarial variables ($\underline{x}_{J_1}$), environmental or chance variables ($\underline{x}_{J_2}$), controlled variables ($\underline{x}_{J_3}$) and query variables ($\underline{x}_{J_0}$). For the inference problem above, each query $\underline{x}_{J_0}$ seeks to maximize the expected minimum payoff of all agents, without observing the adversarial or chance variables, and assuming the adversary makes its decision after observing control and chance variables.

Example 2.8. A “probabilistic” graphical model is defined using a expansion semigroup $\mathcal{G}_e = (\mathbb{R}^{\geq 0}, \times)$ and often a marginalization semigroup $\mathcal{G}_m = (\mathbb{R}^{\geq 0}, +)$. The expanded form represents the unnormalized joint probability $q(\underline{x}) = \prod_I f_I(\underline{x}_I)$, whose marginal probabilities are simply called marginals. Replacing the summation with the

marginalization semigroup $\mathcal{G}_m = (\mathbb{R}^{\geq 0}, \max)$, seeks the maximum probability state and the resulting integration problem $q(\emptyset) = \max_{\underline{x}} \prod_I f_I(\underline{x}_I)$ is known as **maximum a posteriori (MAP)** inference. Alternatively by adding a second marginalization operation to the summation, we get the **marginal MAP** inference

$$q(\underline{x}_{J_0}) = \max_{\underline{x}_{J_2}} \sum_{\underline{x}_{J_1}} \prod_I f_I(\underline{x}_I). \quad (4)$$

where here $\otimes = \prod$, $\oplus^1 = \sum$ and $\oplus^2 = \max$ (recall the numbering of operations is left to right.)

If the object of interest is the negative log-probability (*a.k.a.* energy), the product expansion semigroup is replaced by $\mathcal{G}_e = (\mathbb{R}, +)$. Instead of the sum marginalization semigroup, we can use the **log-sum-exp** semigroup, $\mathcal{G}_m = (\mathbb{R}, +)$ where $a \oplus b \stackrel{\text{def}}{=} \log(e^{-a} + e^{-b})$. The integral in this case is the log-partition function. If we change the marginalization semigroup to $\mathcal{G}_m = (\mathbb{R}, \min)$, the integral is the minimum energy (corresponding to MAP).

3. THE INFERENCE HIERARCHY

Often, the complexity class is concerned with the **decision version** of the inference problem in definition 2.6. The decision version of an inference problem asks a yes/no

question about the integral such as $q(\emptyset) \stackrel{?}{\geq} q^3$ or $q(\emptyset) \stackrel{?}{=} q$ for a given $q \in \mathcal{Y}^*$.

Here, we produce a hierarchy of inference problems in analogy to the polynomial [Stockmeyer 1976], the counting [Wagner 1986] and the arithmetic [Rogers 1987] hierarchies.

To define the hierarchy, we assume the following in definition 2.6:

- Any two consecutive marginalization operations are distinct ($\oplus^l \neq \oplus^{l+1} \forall 1 \leq l < M$).
- The marginalization index sets $J_l \forall 1 \leq l \leq M$ are non-empty. Moreover if $|J_l| = \mathcal{O}(\log(N))$ we call this marginalization operation a **polynomial marginalization** as here $|\mathcal{X}_{J_l}| = \text{Poly}(N)$.
- In defining the factor-graph, we required each factor to be polynomially computable. In building the hierarchy, we require the operations over each semigroup to be polynomially computable as well. To this end we consider the set of rational numbers $\mathcal{Y}^* \subseteq \mathbb{Q}^{\geq 0} \cup \{\pm\infty\}$. Note that this automatically eliminates semigroups that involve operations such as exponentiation and logarithm (because $\mathbb{Q}$ is not closed under these operations) and only consider summation, multiplication, minimization and maximization.

We can always re-express any inference problem to enforce the first two conditions and therefore they do not impose any restriction. Below, we will use a **language** to identify inference problems for an arbitrary set of factors $\mathcal{F} = \{f_I\}$. For example, sum-

product refers to the inference problem $\sum_{\underline{x}} \prod_I f_I(\underline{x}_I) \stackrel{?}{\geq} q$. In this sense the rightmost “token” in the language (here *product*) identifies the expansion semigroup $\mathcal{G}_e = (\mathbb{Q}, \times)$ and the rest of tokens identify the marginalization semigroups over $\mathbb{Q}$ in the given order. Therefore, this minimal language exactly identifies the inference problem. The only information that affects the computational complexity of an inference problem but is not specified in this language is whether each of the marginalization operations are polynomial or exponential.

³Assuming an ordering on $\mathcal{Y}^*$.

We define five **inference families**: $\Sigma, \Pi, \Phi, \Psi, \Delta$. The families are associated with that “outermost” marginalization operation – i.e., $\bigoplus^M$ in definition 2.6. Σ is the family of inference problems where $\bigoplus^M = \text{sum}$. Similarly, Π is associated with product, Φ with minimization and Ψ with maximization. Δ is the family of inference problems where the last marginalization is polynomial (i.e., $|J_M| = \mathcal{O}(\log(N))$ regardless of $\bigoplus^M$).

Now we define **inference classes** in each family, such that all the problems in the same class have the same computational complexity. Here, the hierarchy is exhaustive – i.e., it includes all inference problems using any combination of the four operations sum, min, max and product whenever the integral $q(\emptyset)$ has a polynomial representation (see claim 3.3). Moreover the inference classes are disjoint. For this, each family is parameterized by a subscript M and two sets S and $\mathcal{D}$ (e.g., $\Phi_M(S, \mathcal{D})$ is an inference “class” in family Φ). As before, M is the number of marginalization operations, S is the set of indices of the (exponential) sum-marginalization and $\mathcal{D}$ is the set of indices of polynomial marginalizations.

Example 3.1. Sum-min-sum-product identifies the decision problem

$$\sum_{\underline{x}_{J_3}} \min_{\underline{x}_{J_2}} \sum_{\underline{x}_{J_1}} \prod_I f_I(\underline{x}_I) \stackrel{?}{\geq} q$$

where J_1, J_2 and J_3 partition $\mathcal{N}$. Assume $J_1 = \{2, \dots, \frac{N}{2}\}$, $J_2 = \{\frac{N}{2} + 1, \dots, N\}$ and $J_3 = \{1\}$. Since we have three marginalization operations $M = 3$. Here the first and second marginalizations are exponential and the third one is polynomial (since $|J_3|$ is constant). Therefore $\mathcal{D} = \{3\}$. Since the only exponential summation is $\bigoplus_{\underline{x}_{J_1}}^1 = \sum_{\underline{x}_{J_1}}$, $S = \{1\}$. In our inference hierarchy, this problem belongs to the class $\Delta_3(\{1\}, \{3\})$.

Alternatively, if we use different values for J_1, J_2 and J_3 whose sizes each grow linearly with N , the corresponding inference problem becomes a member of $\Sigma_3(\{1, 3\}, \emptyset)$.

Remark 3.2. Note that arbitrary assignments to M, S and $\mathcal{D}$ do not necessarily define a valid inference class. For example we require that $S \cap \mathcal{D} = \emptyset$ and no index in $\mathcal{D}$ or S can be larger than M . Moreover, the values in S and $\mathcal{D}$ should be compatible with the inference class. For example, for inference class $\Sigma_M(S, \mathcal{D})$, M is a member of S . For notational convenience, if an inference class notation is invalid we equate it with an empty set – e.g., $\Psi_1(\{1\}, \emptyset) = \emptyset$, because $S = \{1\}$ and $M = 1$ means the inference class is Σ rather than Ψ .

In the definition below, we ignore the inference problems in which product appears in any of the marginalization semigroups (e.g., product-sum). The following claim, explains this choice.

CLAIM 3.3. *For $\bigoplus_M = \text{prod}$, the inference query $q(\underline{x}_{J_0})$ can have an exponential representation in N .*

PROOF. The claim states that when the product appears in the marginalization operations, the marginal (and integral) can become very large, such that we can no longer represent them in space polynomial in N . We show this for an integration problem. The same idea can show the exponential representation of a marginal query.

To see why this integral has an exponential representation in N , consider its simplified form

$$q(\emptyset) = \prod_{\underline{x}_I} q(\underline{x}_I)$$

where $q(\underline{x})$ here is the result of inference up to the last marginalization step $\oplus^M$, which is product, where $\mathcal{X}_1$ grows exponentially with N . Recall that the hierarchy is defined for operations on $\mathbb{Q}^{\geq 0}$. Since $q(\underline{x}_1)$ for each $\underline{x}_1 \in \mathcal{X}_1$ has a constant size, say c , the size of representation of $q(\emptyset)$ using a binary scheme is

$$\lceil \log_2(q(\emptyset)) \rceil = \left\lceil \log_2 \left(\prod_{\underline{x}_1} q(\underline{x}_1) \right) \right\rceil = \left\lceil \sum_{\underline{x}_1} c \right\rceil = \lceil c|\mathcal{X}_1| \rceil$$

which is exponential in N . $\square$

Define the **base members** of the families as

$$\begin{aligned} \Sigma_0(\emptyset, \emptyset) &\stackrel{\text{def}}{=} \{\text{sum}\} & \Phi_0(\emptyset, \emptyset) &\stackrel{\text{def}}{=} \{\text{min}\} \\ \Psi_0(\emptyset, \emptyset) &\stackrel{\text{def}}{=} \{\text{max}\} & \Pi_0(\emptyset, \emptyset) &\stackrel{\text{def}}{=} \{\text{prod}\} \\ \Delta_0(\emptyset, \emptyset) &= \emptyset & \Delta_1(\emptyset, \{1\}) &\stackrel{\text{def}}{=} \{\text{sum} - \text{sum}, \text{min} - \text{min}, \text{max} - \text{max}\} \end{aligned} \quad (5)$$

where the initial members of each family only identify the expansion semigroup – e.g., sum in $\Sigma_0(\emptyset, \emptyset)$ identifies $q(\underline{x}) = \sum_{\underline{x}_1} f_1(\underline{x}_1)$. Here, the exception is $\Delta_1(\emptyset, \{1\})$, which contains three *inference problems*.⁴

Let $\Xi_M(\mathcal{S}, \mathcal{D})$ denote the union of corresponding classes within all families:

$$\Xi_M(\mathcal{S}, \mathcal{D}) = \Sigma_M(\mathcal{S}, \mathcal{D}) \cup \Pi_M(\mathcal{S}, \mathcal{D}) \cup \Phi_M(\mathcal{S}, \mathcal{D}) \cup \Psi_M(\mathcal{S}, \mathcal{D}) \cup \Delta_M(\mathcal{S}, \mathcal{D})$$

Now define the **inference family members** recursively, by adding a marginalization operation to all the problems in each inference class. If this marginalization is polynomial then the new class belongs to the Δ family and the set $\mathcal{D}$ is updated accordingly. Alternatively, if this outermost marginalization is exponential, depending on the new marginal operation (i.e., min, max, sum) the new class is defined to be a member of Φ , Ψ or Σ . For the case that the last marginalization is summation set $\mathcal{S}$ is updated.

- **Adding an exponential marginalization** $\forall |\mathcal{X}_{J_M}| = \text{Poly}(N), M > 0$

$$\begin{aligned} \Sigma_{M+1}(\mathcal{S} \cup \{M+1\}, \mathcal{D}) &\stackrel{\text{def}}{=} \{\text{sum} - \xi \mid \xi \in \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Sigma_M(\mathcal{S}, \mathcal{D})\} \\ \Phi_{M+1}(\mathcal{S}, \mathcal{D}) &\stackrel{\text{def}}{=} \{\text{min} - \xi \mid \xi \in \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D})\} \\ \Psi_{M+1}(\mathcal{S}, \mathcal{D}) &\stackrel{\text{def}}{=} \{\text{max} - \xi \mid \xi \in \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Psi_M(\mathcal{S}, \mathcal{D})\} \\ \Pi_{M+1}(\mathcal{S}, \mathcal{D}) &\stackrel{\text{def}}{=} \emptyset \end{aligned} \quad (6)$$

- **Adding a polynomial marginalization** $\forall |\mathcal{X}_{J_M}| = \text{Poly}(N), > 1$

$$\Delta_{M+1}(\mathcal{S}, \mathcal{D} \cup \{M+1\}) \stackrel{\text{def}}{=} \{\oplus - \xi \mid \xi \in \Xi_M(\mathcal{S}, \mathcal{D}), \oplus \in \{\text{min}, \text{max}, \text{sum}\}\} \quad (7)$$

⁴We treat $M = 1$ for Δ specially as in this case the marginalization operation can not be polynomial. This is because if $|J_1| = \mathcal{O}(\log(N))$, then $|J_0| = \Omega(N)$ which violates the conditions in the definition of the inference problem.

3.1. Single marginalization

Following the recursive construction of the hierarchy as defined above, the inference classes in the hierarchy with *one* marginalization are

$$\Delta_1(\emptyset, \{1\}) = \{\min - \min, \max - \max, \text{sum} - \text{sum}\} \quad (8)$$

$$\Psi_1(\emptyset, \emptyset) = \{\max - \min, \max - \text{sum}, \max - \text{prod}\} \quad (9)$$

$$\Phi_1(\emptyset, \emptyset) = \{\min - \max, \min - \text{sum}, \min - \text{prod}\} \quad (10)$$

$$\Sigma_1(\{1\}, \emptyset) = \{\text{sum} - \text{prod}, \text{sum} - \min, \text{sum} - \max\} \quad (11)$$

Now we review all the problems above and prove that Δ_1 , Ψ_1 , Φ_1 and Σ_1 are complete w.r.t. $\mathbb{P}$, NP , coNP and $\mathbb{P}$ respectively. Starting from Δ_1 :

PROPOSITION 3.4. *sum-sum, min-min and max-max inference are in $\mathbb{P}$.*

PROOF. To show that these inference problems are in $\mathbb{P}$, we provide polynomial-time algorithms for them:

- sum – sum is short for

$$q(\emptyset) = \sum_{\underline{x}} \sum_I f_I(\underline{x}_I)$$

which asks for the sum over all assignments of $\underline{x} \in \mathcal{X}$, of the sum of all the factors. It is easy to see that each factor value $f_I(\underline{x}_I) \forall I$, $\mathcal{X}_I$ is counted $|\mathcal{X}_{\setminus I}|$ times in the summation above. Therefore we can rewrite the integral above as

$$q(\emptyset) = \sum_I |\mathcal{X}_{\setminus I}| \left(\sum_{\underline{x}_I} f_I(\underline{x}_I) \right)$$

where the new form involves polynomial number of terms and therefore is easy to calculate.

- min – min (similar for max – max) is short for

$$q(\emptyset) = \min_{\underline{x}} \min_I f_I(\underline{x}_I)$$

where the query seeks the minimum achievable value of any factor. We can easily obtain this by seeking the range of all factors and reporting the minimum value in polynomial time. $\square$

Max-sum and max-prod are widely studied and it is known that their decision version are NP -complete [Shimony 1994]. By reduction from satisfiability we can show that max-min inference [Ravanbakhsh et al. 2014] is also NP -hard.

PROPOSITION 3.5. *The decision version of max-min inference that asks $\max_{\underline{x}} \min_I f_I(\underline{x}_I) \stackrel{?}{\geq} q$ is NP -complete.*

PROOF. Given $\underline{x}$ it is easy to verify the decision problem, so max-min decision belongs to NP . To show NP -completeness, we reduce the 3-SAT to a max-min inference problem, such that 3-SAT is satisfiable *iff* the max-min value is $q(\emptyset) \geq 1$ and unsatisfiable otherwise.

Simply define one factor per clause of 3-SAT, such that $f_I(\underline{x}_I) = 1$ if $\underline{x}_I$ satisfies the clause and any number less than one otherwise. With this construction, the max-min value $\max_{\underline{x}} \min_{I \in \mathcal{F}} f_I(\underline{x}_I)$ is one *iff* the original SAT problem was satisfiable, otherwise it is less than one. This reduces 3-SAT to Max-Min-decision. $\square$

This means all the problems in $\Psi_1(\emptyset, \emptyset)$ are in NP (and in fact are complete w.r.t. this complexity class). In contrast, problems in $\Phi_1(\emptyset, \emptyset)$ are in coNP , which is the class of

decision problems in which the “NO instances” result has a polynomial time verifiable witness or proof. Note that by changing the decision problem from $q(\emptyset) \stackrel{?}{\geq} q$ to $q(\emptyset) \stackrel{?}{\leq} q$, the complexity classes of problems in Φ and Ψ family are reversed (*i.e.*, problems in $\Phi_1(\emptyset, \emptyset)$ become NP -complete and the problems in $\Psi_1(\emptyset, \emptyset)$ become coNP -complete).

Among the members of $\Sigma_1(\{1\}, \emptyset)$, sum-product is known to be PP -complete [Littman et al. 2001; Roth 1993]. It is easy to show the same result for sum-min (sum-max) inference.

PROPOSITION 3.6. *The sum-min decision problem $\sum_{\underline{x}} \min_I f_I(x_I) \stackrel{?}{\geq} q$ is PP -complete for $\mathcal{Y} = \{0, 1\}$.*

PP is the class of problems that are polynomially solvable using a non-deterministic Turing machine, where the acceptance condition is that the majority of computation paths accept.

PROOF. To see that $\sum_{\underline{x}} \min_I f_I(x_I) \stackrel{?}{\geq} q$ is in PP , enumerate all $\underline{x} \in \mathcal{X}$ non-deterministically and for each assignment calculate $\min_I f_I(x_I)$ in polynomial time (where each path accepts iff $\min_I f_I(x_I) = 1$) and accept iff at least q of the paths accept.

Given a matrix $\mathbf{A} \in \{0, 1\}^{N \times N}$ the problem of calculating its permanent

$$\text{perm}(\mathbf{A}) = \sum_{\underline{z} \in \mathcal{S}_N} \prod_{i=1}^N \mathbf{A}_{i, z_i}$$

where $\mathcal{S}_N$ is the set of permutations of $1, \dots, N$ is $\#\text{P}$ -complete and the corresponding decision problem is PP -complete [Valiant 1979]. To show completeness w.r.t. PP it is enough to reduce the problem of computing the matrix permanent to sum-min inference in a graphical model.

The problem of computing the permanent has been reduced to sum-product inference in graphical models [Huang and Jebara 2009]. However, when $f_I(x_I) \in \{0, 1\} \forall I$, sum-product is isomorphic to sum-min. This is because $y_1 \times y_2 = \min(y_1, y_2) \forall y_i \in \{0, 1\}$. Therefore, the problem of computing the permanent for such matrices reduces to sum-min inference in the factor-graph of [Huang and Jebara 2009]. $\square$

3.2. Complexity of general inference classes

Let $\mathcal{U}(\cdot)$ denote the complexity class of an inference class in the hierarchy. In obtaining the complexity class of problems with $M > 1$, we use the following fact, which is also used in the polynomial hierarchy: $\text{P}^{\text{NP}} = \text{P}^{\text{coNP}}$ [Arora and Barak 2009]. In fact $\text{P}^{\text{NP}^{\mathbf{A}}} = \text{P}^{\text{coNP}^{\mathbf{A}}}$, for any oracle $\mathbf{A}$. This means that by adding a polynomial marginalization to the problems in $\Phi_M(\mathcal{S}, \mathcal{D})$ and $\Psi_M(\mathcal{S}, \mathcal{D})$, we get the same complexity class $\mathcal{U}(\Delta_{M+1}(\mathcal{S}, \mathcal{D} \cup \{M+1\}))$. The following gives a recursive definition of complexity class for problems in the inference hierarchy.⁵ Note that the definition of the complexity for each class is very similar to the recursive definition of members of each class in eq (6) and eq (7)

⁵We do not prove the completeness w.r.t. complexity classes beyond the first level of the hierarchy and only assert the membership.

THEOREM 3.7. *The complexity of inference classes in the hierarchy is given by the recursion*

$$\mathcal{U}(\Phi_{M+1}(\mathcal{S}, \mathcal{D})) = \text{coNP}^{\mathcal{U}(\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D}))} \quad (12)$$

$$\mathcal{U}(\Psi_{M+1}(\mathcal{S}, \mathcal{D})) = \text{NP}^{\mathcal{U}(\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Psi_M(\mathcal{S}, \mathcal{D}))} \quad (13)$$

$$\mathcal{U}(\Sigma_{M+1}(\mathcal{S} \cup \{M+1\}, \mathcal{D})) = \text{PP}^{\mathcal{U}(\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Sigma_M(\mathcal{S}, \mathcal{D}))} \quad (14)$$

$$\mathcal{U}(\Delta_{M+1}(\mathcal{S}, \mathcal{D} \cup \{M+1\})) = \text{P}^{\mathcal{U}(\Xi_M(\mathcal{S}, \mathcal{D}))} \quad (15)$$

where the base members are defined in eq (5) and belong to $\mathbb{P}$.

PROOF. Recall that our definition of factor graph ensures that $q(\underline{x})$ can be evaluated in polynomial time and therefore the base members are in $\mathbb{P}$ (for complexity of base members of Δ see theorem 3.4). We use these classes as the base of our induction and assuming the complexity classes above are correct for M , we show that are correct for $M+1$. We consider all the above statements one by one:

- *Complexity for members of $\Phi_{M+1}(\mathcal{S}, \mathcal{D})$:*

Adding an exponential-sized *min*-marginalization to an inference problem with known complexity $\mathbb{A}$, requires a Turing machine to non-deterministically enumerate $\underline{z}_{J_M} \in \mathcal{X}_{J_M}$ possibilities, then call the $\mathbb{A}$ oracle with the “reduced factor-graph” – in which $\underline{x}_{J_M}$ is clamped to $\underline{z}_{J_M}$ – and reject iff any of the calls to oracle rejects. This means $\mathcal{U}(\Phi_{M+1}(\mathcal{S}, \mathcal{D})) = \text{coNP}^{\mathbb{A}}$.

Here, eq (12) is also making another assumption expressed in the following claim.

CLAIM 3.8. *All inference classes in $\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D})$ have the same complexity $\mathbb{A}$.*

- $M = 0$: the fact that $q(\underline{x})$ can be evaluated in polynomial time means that $\mathbb{A} = \mathbb{P}$.
- $M > 0$: $\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D})$ only contains one inference class – that is exactly only one of the following cases is correct:
 - $M \in \mathcal{S} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D}) = \Sigma_M(\mathcal{S}, \mathcal{D})$
 - $M \in \mathcal{D} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D}) = \Delta_M(\mathcal{S}, \mathcal{D})$
 - $M \notin \mathcal{S} \cup \mathcal{D} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D}) = \Psi_M(\mathcal{S}, \mathcal{D})$.
 (in constructing the hierarchy we assume two consecutive marginalizations are distinct and the current marginalization is a minimization.)
- But if $\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D})$ contains a single class, the inductive hypothesis ensures that all problems in $\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Phi_M(\mathcal{S}, \mathcal{D})$ have the same complexity class $\mathbb{A}$.

This completes the proof of our claim.

- *Complexity for members of $\Psi_{M+1}(\mathcal{S}, \mathcal{D})$:*

Adding an exponential-sized *max*-marginalization to an inference problem with known complexity $\mathbb{A}$, requires a Turing machine to non-deterministically enumerate $\underline{z}_{J_M} \in \mathcal{X}_{J_M}$ possibilities, then call the $\mathbb{A}$ oracle with the reduced factor-graph and accept iff any of the calls to oracle accepts. This means $\mathcal{U}(\Psi_{M+1}(\mathcal{S}, \mathcal{D})) = \text{NP}^{\mathbb{A}}$. Here, an argument similar to that of claim 3.8 ensures that $\Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Psi_M(\mathcal{S}, \mathcal{D})$ in eq (13) contains a single inference class.

- *Complexity for members of $\Sigma_{M+1}(\mathcal{S} \cup \{M+1\}, \mathcal{D})$:*

Adding an exponential-sized *sum*-marginalization to an inference problem with known complexity $\mathbb{A}$, requires a Turing machine to non-deterministically enumerate $\underline{z}_{J_M} \in \mathcal{X}_{J_M}$ possibilities, then call the $\mathbb{A}$ oracle with the reduced factor-graph and accept iff majority of the calls to oracle accepts. This means $\mathcal{U}(\Sigma_{M+1}(\mathcal{S}, \mathcal{D})) = \text{PP}^{\mathbb{A}}$.

- $M = 0$: the fact that $q(\underline{x})$ can be evaluated in polynomial time means that $\mathbb{A} = \mathbb{P}$.
- $M > 0$:

- $M \in \mathcal{D} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Sigma_M(\mathcal{S}, \mathcal{D}) = \Delta_M(\mathcal{S}, \mathcal{D})$.
- $M \notin \mathcal{D} \cup \mathcal{S} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) \setminus \Sigma_M(\mathcal{S}, \mathcal{D}) = \Psi_M(\mathcal{S}, \mathcal{D}) \cup \Phi_M(\mathcal{S}, \mathcal{D})$: despite the fact that $\mathbb{A} = \mathcal{U}(\Psi_M(\mathcal{S}, \mathcal{D}))$ is different from $\mathbb{A}' = \mathcal{U}(\Phi_M(\mathcal{S}, \mathcal{D}))$, since $\mathbb{P}\mathbb{P}$ is closed under complement, which means $\mathbb{P}\mathbb{P}^{\mathbb{A}} = \mathbb{P}\mathbb{P}^{\mathbb{A}'}$ and the recursive definition of complexity eq (14) remains correct.

- *Complexity for members of $\Delta_{M+1}(\mathcal{S}, \mathcal{D} \cup \{M+1\})$:*

Adding a polynomial-sized marginalization to an inference problem with known complexity $\mathbb{A}$, requires a Turing machine to deterministically enumerate $\underline{z}_{J_M} \in \mathcal{X}_{J_M}$ possibilities in polynomial time, and each time call the $\mathbb{A}$ oracle with the reduced factor-graph and accept after some polynomial-time calculation. This means $\mathcal{U}(\Psi_{M+1}(\mathcal{S}, \mathcal{D})) = \mathbb{P}^{\mathbb{A}}$. Here, there are three possibilities:

- $M = 0$: here again $\mathbb{A} = \mathbb{P}$.
- $M \in \mathcal{S} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) = \Sigma_M(\mathcal{S}, \mathcal{D})$.
- $M \in \mathcal{D} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) = \Delta_M(\mathcal{S}, \mathcal{D})$.
- $M \notin \mathcal{D} \cup \mathcal{S} \Rightarrow \Xi_M(\mathcal{S}, \mathcal{D}) = \Psi_M(\mathcal{S}, \mathcal{D}) \cup \Phi_M(\mathcal{S}, \mathcal{D})$, in which case since $\mathbb{P}\mathbb{P}^{\mathbb{N}\mathbb{P}^{\mathbb{B}}} = \mathbb{P}\mathbb{P}^{\text{co}\mathbb{N}\mathbb{P}^{\mathbb{B}}}$, the recursive definition of complexity in eq (15) remains correct.

□

Example 3.9. Consider the marginal-MAP inference of eq (4). The decision version of this problem, $q(\emptyset) \stackrel{?}{\geq} q$, is a member of $\Psi_2(\{1\}, \emptyset)$ which also includes max – sum – min and max – sum – max. The complexity of this class according to eq (13) is $\mathcal{U}(\Psi^2(\{1\}, \emptyset)) = \mathbb{N}\mathbb{P}^{\mathbb{P}\mathbb{P}}$. However, marginal-MAP is also known to be “complete” w.r.t. $\mathbb{N}\mathbb{P}^{\mathbb{P}\mathbb{P}}$ [Park and Darwiche 2004]. Now suppose that the max-marginalization over $\underline{x}_{J_2}$ is polynomial (e.g., $|J_2|$ is constant). Then marginal-MAP belongs to $\Delta_2(\{1\}, \{2\})$ with complexity $\mathbb{P}^{\mathbb{P}\mathbb{P}}$. This is because a Turing machine can enumerate all $\underline{z}_{J_2} \in \mathcal{X}_{J_2}$ in polynomial time and call its $\mathbb{P}\mathbb{P}$ oracle to see if

$$q(\underline{x}_{J_0} \mid \underline{z}_{J_2}) \stackrel{?}{\geq} q$$

where $q(\underline{x}_{J_0} \mid \underline{z}_{J_2}) = \sum_{\underline{x}_{J_2}} \prod_I f_I(\underline{x}_{I \setminus J_2}, \underline{z}_{I \cap J_2})$

and *accept* if any of its calls to oracle accepts, and rejects otherwise. Here, $f_I(\underline{x}_{I \setminus J_2}, \underline{z}_{I \cap J_2})$ is the reduced factor, in which all the variables in $\underline{x}_{J_2}$ are fixed to $\underline{z}_{J_2 \cap I}$.

Here, **Toda’s theorem** [Toda 1991] has an interesting implication w.r.t. the hierarchy. This theorem states that $\mathbb{P}\mathbb{P}$ is as hard as the polynomial hierarchy, which means min – max – min – ... – max inference for an arbitrary, but constant, number of min and max operations appears below the sum-product inference in the inference hierarchy.

3.3. Complexity of the hierarchy

By restricting the domain $\mathcal{V}^*$ to $\{0, 1\}$, min and max become isomorphic to logical AND ($\wedge$) and OR ($\vee$) respectively, where $1 \cong \text{TRUE}$, $0 \cong \text{FALSE}$. By considering the restriction of the inference hierarchy to these two operations, we can express quantified satisfiability (QSAT) as inference in a graphical model, where $\wedge \cong \forall$ and $\vee \cong \exists$. Let each factor $f_I(\underline{x}_I)$ be a disjunction –e.g., $f(x_{i,j,k}) = x_i \vee \neg x_j \vee \neg x_k$. Then we have

$$\forall_{\underline{x}_{J_M}} \exists_{\underline{x}_{J_{M-1}}} \dots \exists_{\underline{x}_{J_2}} \forall_{\underline{x}_{J_1}} \bigwedge_I f_I(\underline{x}_I) \cong \min_{\underline{x}_{J_M}} \max_{\underline{x}_{J_{M-1}}} \dots \max_{\underline{x}_{J_2}} \min_{\underline{x}_{J_1}} \bigwedge_I f_I(\underline{x}_I)$$

By adding the summation operation, we can express the stochastic satisfiability [Littman et al. 2001] and by generalizing the constraints from disjunctions we can represent any quantified constraint problem (QCP) [Bordeaux and Monfroy 2002]. QSAT, stochastic SAT and QCPs are all PSPACE-complete, where PSPACE is the class of problems that can be solved by a (non-deterministic) Turing machine in polynomial space. Therefore if we can show that inference in the inference hierarchy is in PSPACE, it follows that inference hierarchy is in PSPACE-complete as well.

ALGORITHM 1: inference in PSPACE

```

input :  $\bigoplus_{\underline{x}_{J_M}}^M \bigoplus_{\underline{x}_{J_{M-1}}}^{M-1} \dots \bigoplus_{\underline{x}_{J_1}}^1 \bigotimes_I f_I(\underline{x}_I)$ 
output:  $q(\underline{x}_{J_0})$ 
for each  $\underline{z}_{J_0} \in \mathcal{X}_{J_0}$  do // loop over the query domain
  for each  $z_{i_N} \in \mathcal{X}_{i_N}$  do // loop over  $\mathcal{X}_{i_N}$ 
    .
    .
    .
    for each  $z_{i_1} \in \mathcal{X}_{i_1}$  do // loop over  $\mathcal{X}_{i_1}$ 
      |  $q_1(z_{i_1}) := \bigotimes_I f_I(\underline{z}_I)$ ;
    end
     $q_{i_2}(z_{i_2}) := \bigoplus_{x_{i_1}}^{j(i_1)} q_1(x_{i_1})$ 
    .
    .
    .
     $q_N(z_{i_N}) := \bigoplus_{x_{i_{N-1}}}^{j(i_{N-1})} q_{N-1}(x_{i_{N-1}})$ 
  end
   $q(\underline{z}_{J_0}) := \bigoplus_{x_{i_N}}^{j(i_N)} q_N(x_{i_N})$ 
end
  
```

THEOREM 3.10. *The inference hierarchy is PSPACE-complete.*

PROOF. To prove that a problem is PSPACE-complete, we have to show that 1) it is in PSPACE and 2) a PSPACE-complete problem reduces to it. We already saw that QSAT, which is PSPACE-complete, reduces to the inference hierarchy. But it is not difficult to show that inference hierarchy is contained in PSPACE. Let

$$q(\underline{x}_{J_0}) = \bigoplus_{\underline{x}_{J_M}}^M \bigoplus_{\underline{x}_{J_{M-1}}}^{M-1} \dots \bigoplus_{\underline{x}_{J_1}}^1 \bigotimes_I f_I(\underline{x}_I)$$

be any inference problem in the hierarchy. We can simply iterate over all values of $\underline{z} \in \mathcal{X}$ in nested loops or using a recursion. Let $j(i) : \{1, \dots, N\} \rightarrow \{1, \dots, M\}$ be the index of the marginalization that involves x_i – that is $i \in J_{j(i)}$. Moreover let $i_1, \dots, i_N$ be an ordering of variable indices such that $j(i_k) \leq j(i_{k+1})$. Algorithm 1 uses this notation to demonstrate this procedure using nested loops. Note that here we loop

over individual domains $\mathcal{X}_{i_k}$ rather than $\mathcal{X}_{J_m}$ and track only temporary tuples q_{i_k} , so that the space complexity remains polynomial in N .

□

4. POLYNOMIAL-TIME INFERENCE

Our definition of inference was based on an expansion operation $\otimes$ and one or more marginalization operations $\overset{1}{\oplus}, \dots, \overset{M}{\oplus}$. If we assume only a single marginalization operation, polynomial time inference is still not generally possible. However, if we further assume that the expansion operation is distributive over marginalization and the factor-graph has no loops, exact polynomial time inference is possible.

Definition 4.1. A **commutative semiring** $\mathcal{S} = (\mathcal{Y}^*, \oplus, \otimes)$ is the combination of two commutative semigroups $\mathcal{G}_e = (\mathcal{Y}^*, \otimes)$ and $\mathcal{G}_m = (\mathcal{Y}^*, \oplus)$ with two additional properties

- identity elements $\overset{\oplus}{1}$ and $\overset{\otimes}{1}$ such that $\overset{\oplus}{1} \oplus a = a$ and $\overset{\otimes}{1} \otimes a = a$. Moreover $\overset{\oplus}{1}$ is an **annihilator** for $\mathcal{G}_e = (\otimes, \mathcal{Y}^*)$: $a \otimes \overset{\oplus}{1} = \overset{\oplus}{1} \quad \forall a \in \mathcal{Y}^*$.⁶
- distributive property:

$$a \otimes (b \oplus c) = (a \otimes b) \oplus (a \otimes c) \quad \forall a, b, c \in \mathcal{Y}^*$$

The mechanism of efficient inference using distributive law can be seen in a simple example: instead of calculating $\min(a + b, a + c)$, using the fact that summation distributes over minimization, we may instead obtain the same result using $a + \min(b, c)$, which requires fewer operations.

Example 4.2. The following are some examples of commutative semirings:

- Sum-product $(\mathbb{R}^{\geq 0}, +, \times)$.
- Max-product $(\mathbb{R}^{\geq 0} \cup \{-\infty\}, \max, \times)$ and $(\{0, 1\}, \max, \times)$.
- Min-max $(\mathcal{S}, \min, \max)$ on any ordered set $\mathcal{S}$.
- Min-sum $(\mathbb{R} \cup \{\infty\}, \min, +)$ and $(\{0, 1\}, \min, +)$.
- Or-and $(\{\text{TRUE}, \text{FALSE}\}, \vee, \wedge)$.
- Union-intersection $(2^{\mathcal{S}}, \cup, \cap)$ for any power-set $2^{\mathcal{S}}$.
- The semiring of natural numbers with greatest common divisor and least common multiple $(\mathcal{N}, \text{lcm}, \text{gcd})$.
- Symmetric difference-intersection semiring for any power-set $(2^{\mathcal{S}}, \nabla, \cap)$.

Many of the semirings above are isomorphic –e.g., $y' \cong -\log(y)$ defines an isomorphism between min-sum and max-product. It is also easy to show that the or-and semiring is isomorphic to min-sum/max-product semiring on $\mathcal{Y}^* = \{0, 1\}$.

The inference problems in the example above have different properties indirectly inherited from their commutative semirings: for example, the operation $\min$ (also $\max$) is a **choice function**, which means $\min_{a \in \mathcal{A}} a \in \mathcal{A}$. The implication is that if the summation ($\oplus$) of the semiring is $\min$ (or $\max$), we can replace it with $\arg_{\mathcal{S}_{J_M}} \max$ and (if required) recover $q(\emptyset)$ using $q(\emptyset) = \bigotimes_I f_I(x^*)$ in polynomial time.

As another example, since both operations have inverses, sum-product is a **field** [Pinter 2012]. The availability of inverse for $\otimes$ operation – i.e., when $\mathcal{G}_e$ is an Abelian group – has an important implication for inference: the expanded form of eq (1) can be

⁶When dealing with reals, this is $\overset{\oplus}{1} = 0$; note $a \times 0 = 0$. Indeed it may be useful to view $\overset{\otimes}{1}$ as $1 \in \mathbb{R}$ and $\overset{\oplus}{1}$ as $0 \in \mathbb{R}$.

normalized, and we may inquire about **normalized marginals**

$$p(\underline{x}_J) = \bigoplus_{\underline{x}_{\setminus J}} p(\underline{x}) \quad (16)$$

$$\text{where } p(\underline{x}) \stackrel{\text{def}}{=} \frac{1}{q(\emptyset)} \otimes \left(\bigotimes_I f_I(\underline{x}_I) \right) \quad \text{if } q(\emptyset) \neq \overset{\oplus}{1} \quad (17)$$

$$p(\underline{x}) \stackrel{\text{def}}{=} \overset{\oplus}{1} \quad \text{if } q(\emptyset) = \overset{\oplus}{1} \quad (18)$$

where $p(\underline{x})$ is the normalized joint form. We deal with the case where the integral evaluates to the annihilator as a special case because division by annihilator may not be well-defined. This also means, when working with normalized expanded form and normalized marginals, we always have $\bigoplus_{\underline{x}_J} p(\underline{x}_J) = \overset{\otimes}{1}$

Example 4.3. Since $\mathcal{G}_e = (\mathbb{R}^{>0}, \times)$ and $\mathcal{G}_e = (\mathbb{R}, +)$ are both Abelian groups, min-sum and sum-product inference have normalized marginals. For min-sum inference this means $\min_{\underline{x}_J} p(\underline{x}_J) = \overset{\text{sum}}{1} = 0$. However, for min-max inference, since $(\mathcal{S}, \max)$ is not Abelian, normalized marginals are not defined.

We can apply the identity and annihilator of a commutative semiring to define constraints.

Definition 4.4. A **constraint** is a factor $f_I : \mathcal{X}_I \rightarrow \{\overset{\otimes}{1}, \overset{\oplus}{1}\}$ whose range is limited to identity and annihilator of the expansion monoid.⁷

Here, $f_I(\underline{x}) = \overset{\oplus}{1}$ iff $\underline{x}$ is forbidden and $f_I(\underline{x}) = \overset{\otimes}{1}$ iff it is permissible. A **constraint satisfaction problem** (CSP) is any inference problem on a semiring in which all factors are constraints. Note that this allows definition of the “same” CSP on any commutative semiring and also indicates that inference in general semirings should be difficult. The following theorem formalizes this intuition.

THEOREM 4.5. *Inference in any commutative semiring is NP-hard under randomized polynomial-time reduction.*

PROOF. To prove that inference in any semiring $\mathcal{S} = (\mathcal{Y}^*, \overset{\oplus}{1}, \overset{\otimes}{1})$ is NP-hard under randomized polynomial reduction, we deterministically reduce *unique satisfiability* (USAT) to an inference problems on any semiring. USAT is a so-called “promise problem”, that asks whether a satisfiability problem that is promised to have either zero or one satisfying assignment is satisfiable. [Valiant and Vazirani 1986] prove that a polynomial time randomized algorithm (RP) for USAT implies that $\text{RP} = \text{NP}$.

For this reduction consider a set of binary variables $\underline{x} \in \{0, 1\}^N$, one per each variable in the given instance of USAT. For each clause, define a constraint factor f_I such that $f_I(\underline{x}_I) = \overset{\otimes}{1}$ if $\underline{x}_I$ satisfies that clause and $f_I(\underline{x}_I) = \overset{\oplus}{1}$ otherwise. This means, $\underline{x}$ is a satisfying assignment for USAT iff $q(\underline{x}) = \bigotimes_I f_I(\underline{x}_I) = \overset{\otimes}{1}$. If the instance is unsatisfiable, the integral $q(\emptyset) = \bigoplus_{\underline{x}} \overset{\oplus}{1} = \overset{\oplus}{1}$ (by definition of $\overset{\oplus}{1}$). If the instance is satisfiable there is only a single instance $\underline{x}^*$ for which $q(\underline{x}^*) = \overset{\otimes}{1}$, and therefore the integral evaluates to

⁷Recall that a monoid is a semigroup with an identity. The existence of identity here is a property of the semiring.

1. Therefore we can decide the satisfiability of USAT by performing inference on any semiring, by only relying on the properties of identities.

□

Example 4.6. Inference on xor-and semiring $(\{\text{TRUE}, \text{FALSE}\}, \text{xor}, \wedge)$, where each factor has a disjunction form, is called parity-SAT, which asks whether the number of SAT solutions is even or odd. A corollary to theorem 4.5 is that parity-SAT is $\mathbb{NP}$ -hard under randomized reduction, which is indeed the case [Valiant and Vazirani 1986].

We find it useful to use the same notation for the **identity function** $1(\text{condition})$ – e.g.,

$$1(\text{cond.}) \stackrel{\text{def}}{=} \begin{cases} (+, \times) & (\min, +) & (\min, \max) \\ \text{cond.} = \text{TRUE} & 1 & 0 & -\infty \\ \text{cond.} = \text{FALSE} & 0 & +\infty & +\infty \end{cases} \quad (19)$$

where the intended semiring for $1(\cdot)$ function will be clear from the context.

4.1. Distributive law and its limits

A naive approach to inference over commutative semirings

$$q(\underline{x}_J) = \bigoplus_{\underline{x}_{J \setminus I}} \bigotimes_I f_I(\underline{x}_I) \quad (20)$$

or its normalized version (eq (16)), involves constructing a complete N -dimensional array of $q(\underline{x})$ using the tensor product $q(\underline{x}) = \bigotimes_I f_I(\underline{x}_I)$ and then perform $\oplus$ -marginalization. However, the number of elements in $q(\underline{x})$ is $|\mathcal{X}|$, which is exponential in N , the number of variables.

If the factor-graph is loop free, we can use distributive law to make inference tractable. Assuming $q(\underline{x}_K)$ (or $q(x_k)$) is the marginal of interest, form a tree with K (or k) as its root. Then starting from the leaves, using the distributive law, we can move the $\oplus$ inside the $\otimes$ and define “messages” from leaves towards the root as follows:

$$q_{i \rightarrow I}(x_i) = \bigotimes_{J \in \partial i \setminus I} q_{J \rightarrow i}(x_i) \quad (21)$$

$$q_{I \rightarrow i}(x_i) = \bigoplus_{\underline{x}_{I \setminus i}} f_I(\underline{x}_I) \bigotimes_{j \in \partial I \setminus i} q_{j \rightarrow I}(x_j) \quad (22)$$

where eq (21) defines the message from a variable to a factor, closer to the root and similarly eq (22) defines the message from factor I to a variable i closer to the root. Here, the distributive law allows moving the $\oplus$ over the domain $\mathcal{X}_{I \setminus i}$ from outside to inside of eq (22) – the same way $\oplus$ moves its place in $(a \otimes b) \oplus (a \otimes c)$ to give $a \otimes (b \oplus c)$, where a is analogous to a message.

By starting from the leaves, and calculating the messages towards the root, we obtain the marginal over the root node as the product of incoming messages

$$q(x_k) = \bigotimes_{I \in \partial k} q_{I \rightarrow k}(x_k) \quad (23)$$

In fact, we can assume any subset of variables $\underline{x}_A$ (and factors within those variables) to be the root. Then, the set of all incoming messages to A , produces the marginal

$$q(\underline{x}_A) = \left(\bigotimes_{I \subseteq A} f_I(\underline{x}_I) \right) \left(\bigotimes_{i \in A, J \in \partial i, J \not\subseteq A} q_{J \rightarrow i}(\underline{x}_i) \right) \quad (24)$$




Fig. 2. The figure shows a loop-free factor-graph and the direction of messages sent between variable and factor nodes in order to calculate the marginal over the grey region.

Example 4.7. Consider the joint form represented by the factor-graph of figure 2

$$q(\underline{x}) = \bigotimes_{A \in \{I, J, K, L, O, T, U, V, W, X, Y, Z\}} f_A(\underline{x}_A)$$

and the problem of calculating the marginal over $\underline{x}_{\{i, j, k\}}$ (i.e., the shaded region).

$$q(\underline{x}_{\{i, j, k\}}) = \bigoplus_{\underline{x}_{\setminus \{i, j, k\}}} \bigotimes_{A \in \{I, J, K, L, O, T, U, V, W, X, Y, Z\}} f_A(\underline{x}_A)$$

We can move the $\oplus$ inside the $\otimes$ to obtain

$$q(\underline{x}_{\{i, j, k\}}) = f_I(\underline{x}_I) \otimes q_{L \rightarrow i}(x_i) \otimes q_{K \rightarrow i}(x_i) \otimes q_{V \rightarrow j}(x_j) \otimes q_{W \rightarrow j}(x_j) \otimes q_{K \rightarrow k}(x_k)$$

where each term $q_{A \rightarrow i}$ factors the summation on the corresponding sub-tree. For example

$$q_{L \rightarrow i} = \bigoplus_{x_w} f_L(\underline{x}_L)$$

Here the message $q_{W \rightarrow j}$ is itself a computational challenge

$$q_{W \rightarrow j} = \bigoplus_{\underline{x}_{\setminus j}} \bigotimes_{A \in \{W, U, Y, X, O, T, Z\}} f_A(\underline{x}_A)$$

However we can also decompose this message over sub-trees

$$q_{W \rightarrow j} = \bigoplus_{\underline{x}_{\setminus j}} f_A(\underline{x}_A) \otimes q_{e \rightarrow W}(x_e) \otimes q_{r \rightarrow W}(x_r)$$

where again, using the distributive law, $q_{e \rightarrow W}$ and $q_{r \rightarrow W}$ further simplify based on the incoming messages to the variable nodes x_r and x_e .

This procedure is known as Belief Propagation (BP), which is sometimes prefixed with the corresponding semiring – e.g., sum-product BP. Even though BP is only guaranteed to produce correct answers when the factor-graph is a tree (and few other cases [Aji et al. 1998; Weiss 2001; Bayati et al. 2005; Weller and Jebara 2013]), it performs

surprisingly well when applied as a fixed point iteration to graphs with loops [Murphy et al. 1999; Gallager 1962]. In the case of loopy graphs, the message updates are repeatedly applied in the hope of convergence. Here, for numerical stability, when the $\otimes$ operator has an inverse, the messages are normalized. We use $\propto$ to indicate this normalization according to the mode of inference

$$\hat{p}_{I \rightarrow i}(x_i) \propto \bigoplus_{\underline{x}_{\setminus i}} f_I(\underline{x}_I) \bigotimes_{j \in \partial I \setminus i} \hat{p}_{j \rightarrow I}(x_j) \propto P_{I \rightarrow i}(\hat{\underline{p}}_{\partial I \setminus i \rightarrow I})(x_i) \quad (25)$$

$$\hat{p}_{i \rightarrow I}(x_i) \propto \bigotimes_{J \in \partial i \setminus I} \hat{p}_{J \rightarrow i}(x_i) \propto P_{i \rightarrow I}(\hat{\underline{p}}_{\partial i \setminus I \rightarrow i})(x_i) \quad (26)$$

$$\hat{p}(\underline{x}_I) \propto f_I(\underline{x}_I) \bigotimes_{i \in \partial I} \hat{p}_{i \rightarrow I}(x_i) \quad (27)$$

$$\hat{p}(x_i) \propto \bigotimes_{I \in \partial i} \hat{p}_{I \rightarrow i}(x_i) \quad (28)$$

Here, for general graphs, $\hat{p}(x_i)$ and $\hat{p}(\underline{x}_I)$ are approximations to $p(x_i)$ and $p(\underline{x}_I)$ of eq (16). The **functionals** $P_{i \rightarrow I}(\hat{\underline{p}}_{\partial i \setminus I \rightarrow i})(\cdot)$ and $P_{I \rightarrow i}(\hat{\underline{p}}_{\partial I \setminus i \rightarrow I})(\cdot)$ cast the BP message updates as an operator on a subset of incoming messages – i.e., $\hat{\underline{p}}_{\partial i \setminus I \rightarrow i} = \{\hat{p}_{J \rightarrow i} \mid J \in \partial i \setminus I\}$. We use these functional notation in presenting the algebraic form of survey propagation in section 5.

4.2. The limits of message passing

By observing the application of distributive law in semirings, a natural question to ask is: can we use distributive law for polynomial time inference on loop-free graphical models over any of the inference problems at higher levels of inference hierarchy or in general, for any inference problem with more than one marginalization operation? The answer to this question is further motivated by the fact that, when loops exists, the same scheme may become a powerful approximation technique. When we have more than one marginalization operations, a natural assumption in using distributive law is that the expansion operation distributes over all the marginalization operations – e.g., as in min-max-sum (where sum distributes over both min and max), min-max-min, xor-or-and. Consider the simplest case with three operators $\overset{1}{\oplus}$, $\overset{2}{\oplus}$ and $\otimes$, where $\otimes$ distributes over both $\overset{1}{\oplus}$ and $\overset{2}{\oplus}$. Here the integration problem is

$$q(\emptyset) = \bigoplus_{\underline{x}_{J_2}}^2 \bigoplus_{\underline{x}_{J_1}}^1 \bigotimes_I f_I(\underline{x}_I)$$

where J_1 and J_2 partition $\{1, \dots, N\}$.

In order to apply distributive law for each pair $(\overset{1}{\oplus}, \otimes)$ and $(\overset{2}{\oplus}, \otimes)$, we need to be able to commute $\overset{1}{\oplus}$ and $\overset{2}{\oplus}$ operations. That is, we require

$$\bigoplus_{\underline{x}_A}^1 \bigoplus_{\underline{x}_B}^2 g(\underline{x}_{A \cup B}) = \bigoplus_{\underline{x}_B}^2 \bigoplus_{\underline{x}_A}^1 g(\underline{x}_{A \cup B}). \quad (29)$$

for the specified $A \subseteq J_1$ and $B \subseteq J_2$.

Now, consider a simple case involving two binary variables x_i and x_j , where $g(\underline{x}_{\{i,j\}})$ is

$$\begin{array}{c}
 \begin{array}{cc}
 & x_j \\
 & 0 \quad 1 \\
 x_i \begin{array}{c} 0 \\ 1 \end{array} & \begin{array}{|cc|}
 \hline
 a & b \\
 \hline
 c & d \\
 \hline
 \end{array}
 \end{array}
 \end{array}$$

Applying eq (29) to this simple case (*i.e.*, $A = \{i\}, B = \{j\}$), we require

$$(a \oplus^1 b) \oplus^2 (c \oplus^1 d) = (a \oplus^2 b) \oplus^1 (c \oplus^2 d).$$

The following theorem leads immediately to a negative result:

THEOREM 4.8. [Eckmann and Hilton 1962]:

$$(a \oplus^1 b) \oplus^2 (c \oplus^1 d) = (a \oplus^2 b) \oplus^1 (c \oplus^2 d) \Leftrightarrow \oplus^1 = \oplus^2 \quad \forall a, b, c$$

which implies that *direct application of distributive law to tractably and exactly solve any inference problem with more than one marginalization operation is unfeasible, even for tree structures*. This limitation was previously known for marginal MAP inference [Park and Darwiche 2004].

Min and max operations have an interesting property in this regard. Similar to any other operations for min and max we have

$$\min_{x_J} \max_{x_I} g(x_{I \cup J}) \neq \max_{x_I} \min_{x_J} g(x_{I \cup J})$$

However, if we slightly change the inference problem (from pure assignments $x_{J_i} \in \mathcal{X}_{J_i}$ to a distribution over assignments; *a.k.a.* mixed strategies), as a result of the celebrated *minimax theorem* [Von Neumann and Morgenstern 2007], the min and max operations commute – *i.e.*,

$$\min_{s(x_J)} \max_{s(x_I)} \sum_{x_{I \cup J}} s(x_J) g(x_{I \cup J}) s(x_I) = \max_{s(x_I)} \min_{s(x_J)} \sum_{x_{I \cup J}} s(x_I) g(x_{I \cup J}) s(x_J)$$

where $s(x_{J_1})$ and $s(x_{J_2})$ are mixed strategies. This property has enabled addressing problems with min and max marginalization operations using message-passing-like procedures. For example, [Ibrahimi et al. 2011] solve this (mixed-strategy) variation of min-max-product inference. Message passing procedures that operate on graphical models for game theory (*a.k.a.* “graphical games”) also rely on this property [Ortiz and Kearns 2002; Kearns 2007].

5. ALGEBRA OF SURVEY PROPAGATION

Survey propagation (SP) was first introduced as a message passing solution to satisfiability [Braunstein et al. 2002a] and was later generalized to general CSP [Braunstein et al. 2002b] and arbitrary inference problems over factor-graphs [Mézard and Montanari 2009]. Several works offer different interpretations and generalizations of survey propagation [Kroc et al. 2002; Braunstein and Zecchina 2003; Maneva et al. 2004]. Here, we propose a generalization based the same notions that extends the application of BP to arbitrary commutative semirings. Our derivation generalizes the variational approach of [Mézard and Montanari 2009], in the same way that the algebraic approach to BP (using commutative semirings) generalizes the variational derivation of sum-product and min-sum BP.

As a fixed point iteration procedure, if BP has more than one fixed points, it may not converge at all. Alternatively, if the messages are initialized properly, BP may converge to one of its fixed points. SP equations take “all” BP fixed points into account. We view this task, of dealing with all fixed points, as using a third binary operation $\hat{\oplus}$ on $\mathcal{Y}^*$.

In particular, we require that $\otimes$ also distribute over $\hat{\oplus}$, forming a second commutative semiring. We refer to this new semiring as a **SP semiring**. To better explain the role of the third operation we need to introduce some notation.

Let $\hat{\underline{p}}_{\rightarrow.}$ be a BP fixed point – that is let

$$\hat{\underline{p}}_{\rightarrow.} = \{ \hat{p}_{i \rightarrow I} = P_{i \rightarrow I}(\hat{\underline{p}}_{\partial i \setminus I \rightarrow i}), \hat{p}_{I \rightarrow i} = P_{I \rightarrow i}(\hat{\underline{p}}_{\partial I \setminus i \rightarrow I}) \mid \forall i, I \in \partial i \}$$

and denote the set of all such fixed points by $\mathcal{P}$. Each BP fixed point corresponds to an approximation to the integral $q(\emptyset)$, which we denote by $Q(\hat{\underline{p}}_{\rightarrow.})(\emptyset)$ – using this functional form to emphasize the dependence of this approximation on BP messages. Recall that in the original problem, $\mathcal{X}$ is the domain of assignments, $q(\underline{x})$ is the expanded form and $\oplus$ -marginalization is (approximately) performed by BP. In the case of survey propagation, $\mathcal{P}$ is the domain of assignments and the integral $Q(\hat{\underline{p}}_{\rightarrow.})(\emptyset)$ evaluates a particular assignment $\hat{\underline{p}}_{\rightarrow.}$ to all the messages – i.e., $Q(\hat{\underline{p}}_{\rightarrow.})(\emptyset)$ is the new expanded form.

In this algebraic perspective, SP efficiently performs a second integral using $\hat{\oplus}$ over all fixed points:

$$Q(\emptyset)(\emptyset) = \hat{\oplus}_{\hat{\underline{p}}_{\rightarrow.} \in \mathcal{P}} Q(\hat{\underline{p}}_{\rightarrow.})(\emptyset) \quad (30)$$

Table I summarizes this correspondence. It is not immediately obvious how this second integration can be useful in practice. The following examples attempt to motivate SP before we explain its derivation.

Example 5.1. If the $\otimes$ operator of the semiring has an inverse, any BP fixed point $\hat{\underline{p}}_{\rightarrow.}$ represents a joint form $\hat{p}(\underline{x})$. We can explicitly write this distribution based on BP marginals (ergo BP messages):

$$\hat{p}(\underline{x}) = \frac{\otimes_I \hat{p}(x_I)}{\otimes_i (\hat{p}(x_i) \odot (|\partial i| - 1))} = \hat{P}(\hat{\underline{p}}_{\rightarrow.})(\underline{x}) \quad (31)$$

where the inverse is w.r.t $\otimes$ and the *exponentiation operator* is defined as $a \odot b \stackrel{\text{def}}{=} \underbrace{a \otimes \dots \otimes a}_{b \text{ times}}$. Our notation using $\hat{P}(\hat{\underline{p}}_{\rightarrow.})(\underline{x})$ is to explicitly show the dependence of this

joint form on BP messages. To see why this is correct, we use the exactness of BP on trees and substitute BP marginals eqs (28) and (27) into eq (31):

$$\begin{aligned} \frac{\otimes_I \hat{p}(x_I)}{\otimes_i (\hat{p}(x_i) \odot (|\partial i| - 1))} &= \frac{\otimes_I f_I(\underline{x}_I) \otimes_{i \in \partial I} \hat{p}_{i \rightarrow I}(x_i)}{\otimes_i (\otimes_{I \in \partial i} \hat{p}_{I \rightarrow i}(x_i) \odot (|\partial i| - 1))} = \\ \frac{\otimes_I f_I(\underline{x}_I) \otimes_{i \in \partial I} \hat{p}_{i \rightarrow I}(x_i)}{\otimes_i (\otimes_{I \in \partial i} \hat{p}_{I \rightarrow i}(x_i))} &= \otimes_I f_I(\underline{x}_I) = p(\underline{x}) \end{aligned}$$

In some interesting settings for sum-product inference, BP has many “disjoint” fixed points, in the sense that $\hat{p}(\underline{x})$ estimates (as defined above) have small overlap with each other (e.g., citepMezard1987) – i.e., $p(\underline{x}) \approx \sum_{\hat{\underline{p}}_{\rightarrow.} \in \mathcal{P}} \hat{P}(\hat{\underline{p}}_{\rightarrow.})(\underline{x})$. In this setting the BP estimate of the integral (and marginals) is inaccurate. However, summing over all such BP integrals gives a more accurate estimate of $q(\emptyset)$. This is an application of SP where $\oplus = \hat{\oplus} = +$ in eq (30).

Table I. The correspondance between BP and SP

Belief Propagation	Survey Propagation
domain:	
$\forall i \quad \begin{array}{c} \underline{x} \\ x_i \\ \mathcal{X} \end{array}$	$\begin{array}{c} \widehat{\underline{p}}_{\rightarrow} \\ \widehat{\underline{p}}_{i \rightarrow I}, \widehat{\underline{p}}_{I \rightarrow i} \\ \mathcal{P} \end{array} \quad \forall i, I \in \partial i$
expanded form:	
$q(\underline{x})$	$Q(\widehat{\underline{p}}_{\rightarrow})(\emptyset)$
integration:	
$q(\emptyset) = \bigoplus_{\underline{x}} q(\underline{x})$	$Q(\emptyset) = \bigoplus_{\widehat{\underline{p}}_{\rightarrow}} Q(\widehat{\underline{p}}_{\rightarrow})(\emptyset)$
marginalization:	
$p(x_i) \propto \bigoplus_{\underline{x} \setminus i} p(\underline{x})$	$S(\widehat{\underline{p}}_{I \rightarrow i}) \propto \bigoplus_{\widehat{\underline{p}}_{I \rightarrow i}} P(\widehat{\underline{p}}_{\rightarrow})$
factors:	
$\forall I \quad f_I(\underline{x}_I)$	$\tilde{P}_I(\widehat{\underline{p}}_{\partial I \rightarrow I})(\emptyset), \tilde{P}_i(\widehat{\underline{p}}_{\partial i \setminus I \rightarrow i})(\emptyset) \text{ and } \tilde{P}_{i \leftrightarrow I}(\widehat{\underline{p}}_{i \rightarrow I}, \widehat{\underline{p}}_{I \rightarrow i})(\emptyset)^{-1} \quad \forall i, I \in \partial i$

Our derivation of SP requires $(\mathcal{Y}^*, \otimes)$ to be an Abelian group (*i.e.*, every element of $\mathcal{Y}^*$ has an inverse w.r.t. $\otimes$). We require $\otimes$ to be invertable as we need to work with normalized BP and SP messages. In section 5.4 we introduce another variation of SP that simply counts the BP fixed points and relaxes this requirement.

5.1. Decomposition of the integral

In writing the normalized BP equations in section 4.1, we hid the normalization constant using $\propto$ sign. Here we explicitly define the normalization constants or **local integrals** by defining unnormalized messages, based on their normalized version

$$\tilde{p}_{I \rightarrow i}(x_i) \stackrel{\text{def}}{=} \bigoplus_{\underline{x} \setminus i} f_I(\underline{x}_I) \bigotimes_{j \in \partial I \setminus i} \widehat{p}_{j \rightarrow I}(x_j) \stackrel{\text{def}}{=} \tilde{P}_{I \rightarrow i}(\widehat{\underline{p}}_{\partial I \setminus i \rightarrow I})(x_i) \quad (32)$$

$$\tilde{p}_{i \rightarrow I}(x_i) \stackrel{\text{def}}{=} \bigotimes_{J \in \partial i \setminus I} \widehat{p}_{J \rightarrow i}(x_i) \stackrel{\text{def}}{=} \tilde{P}_{i \rightarrow I}(\widehat{\underline{p}}_{\partial i \setminus I \rightarrow i})(x_i) \quad (33)$$

$$\tilde{p}_I(\underline{x}_I) \stackrel{\text{def}}{=} f_I(\underline{x}_I) \bigotimes_{i \in \partial I} \widehat{p}_{i \rightarrow I}(x_i) \stackrel{\text{def}}{=} \tilde{P}_I(\widehat{\underline{p}}_{\partial I \rightarrow I})(\underline{x}_I) \quad (34)$$

$$\tilde{p}_i(x_i) \stackrel{\text{def}}{=} \bigotimes_{I \in \partial i} \widehat{p}_{I \rightarrow i}(x_i) \stackrel{\text{def}}{=} \tilde{P}_i(\widehat{\underline{p}}_{\partial i \rightarrow i})(x_i) \quad (35)$$

where each update also has a functional form on the r.h.s. In each case, the local integrals are simply the integral of unnormalized messages or marginals – *e.g.*, $\tilde{p}_{I \rightarrow i}(\emptyset) = \bigoplus_{x_i} \tilde{P}_{I \rightarrow i}(x_i)$.

Define the functional $\tilde{P}_{i \leftrightarrow I}(\widehat{\underline{p}}_{i \rightarrow I}, \widehat{\underline{p}}_{I \rightarrow i})$ as the product of messages from i to I and vice versa

$$\tilde{P}_{i \leftrightarrow I}(x_i) \stackrel{\text{def}}{=} \widehat{p}_{i \rightarrow I}(x_i) \otimes \widehat{p}_{I \rightarrow i}(x_i) \stackrel{\text{def}}{=} \tilde{P}_{i \leftrightarrow I}(\widehat{\underline{p}}_{i \rightarrow I}, \widehat{\underline{p}}_{I \rightarrow i})(x_i) \quad (36)$$

THEOREM 5.2. *If the factor-graph has no loops and $(\mathcal{Y}^*, \otimes)$ is an Abelian group, the global integral decomposes to local BP integrals as*

$$q(\emptyset) = \bigotimes_I \tilde{p}_I(\emptyset) \bigotimes_i \tilde{p}_i(\emptyset) \left(\bigotimes_{i, I \in \partial i} \tilde{P}_{i \leftrightarrow I}(\emptyset) \right)^{-1} \quad (37)$$

or in other words $q(\emptyset) = Q(\hat{\underline{p}}_{\rightarrow})(\emptyset)$ where

$$Q(\hat{\underline{p}}_{\rightarrow})(\emptyset) = \bigotimes_I \tilde{P}_I(\hat{\underline{p}}_{\partial I \rightarrow I})(\emptyset) \bigotimes_i \tilde{P}_i(\hat{\underline{p}}_{\partial i \rightarrow i})(\emptyset) \left(\bigotimes_{i, I \in \partial i} \tilde{P}_{i \rightarrow I}(\hat{\underline{p}}_{i \rightarrow I}, \hat{\underline{p}}_{I \rightarrow i})(\emptyset) \right)^{-1} \quad (38)$$

PROOF. For this proof we build a tree around an root node r that is connected to one factor. (Since the factor-graph is a tree such a node always exists.) Send BP messages from the leaves, up towards the root r and back to the leaves. Here, any message $q_{i \rightarrow I}(x_i)$, can give us the integral for the sub-tree that contains all the nodes and factors up to node i using $q_{i \rightarrow I}(\emptyset) = \bigoplus_{x_i} q_{i \rightarrow I}(x_i)$. Noting that the root is connected to exactly one factor, the global integral is

$$\bigoplus_{x_r} q(x_r) = \bigoplus_{x_r} \bigotimes_{I \in \partial r} q_{I \rightarrow r}(x_r) = q_{I \rightarrow r}(\emptyset) \quad (39)$$

On the other hand, we have the following relation between $q_{i \rightarrow I}$ and $\hat{\underline{p}}_{i \rightarrow I}$ (also corresponding factor-to-variable message)

$$q_{i \rightarrow I}(x_i) = \hat{\underline{p}}_{i \rightarrow I}(x_i) \otimes q_{i \rightarrow I}(\emptyset) \quad \forall i, I \in \partial i \quad (40)$$

$$q_{I \rightarrow i}(x_i) = \hat{\underline{p}}_{I \rightarrow i}(x_i) \otimes q_{I \rightarrow i}(\emptyset) \quad \forall i, I \in \partial i \quad (41)$$

Substituting this into BP eqs (21) and (22) we get

$$q_{i \rightarrow I}(x_i) = \bigotimes_{J \in \partial i \setminus I} q_{J \rightarrow i}(\emptyset) \hat{\underline{p}}_{J \rightarrow i}(x_i) \quad (42)$$

$$q_{I \rightarrow i}(x_i) = \bigoplus_{\underline{x}_{\setminus i}} f_I(\underline{x}_I) \bigotimes_{j \in \partial I \setminus i} q_{j \rightarrow I}(\emptyset) \hat{\underline{p}}_{j \rightarrow I}(x_j) \quad (43)$$

By summing over both l.h.s and r.h.s in equations above and substituting from eq (33), we get

$$\begin{aligned} \bigoplus_{x_i} q_{i \rightarrow I}(x_i) &= \left(\bigotimes_{J \in \partial i \setminus I} q_{J \rightarrow i}(\emptyset) \right) \otimes \left(\bigoplus_{x_i} \bigotimes_{J \in \partial i \setminus I} \hat{\underline{p}}_{J \rightarrow i}(x_i) \right) \Rightarrow \\ q_{i \rightarrow I}(\emptyset) &= \tilde{\underline{p}}_{i \rightarrow I}(\emptyset) \bigotimes_{J \in \partial i \setminus I} q_{J \rightarrow i}(\emptyset) \end{aligned} \quad (44)$$

and similarly for eq (43) using integration and substitution from eq (32) we have

$$\begin{aligned} \bigoplus_{x_i} q_{I \rightarrow i}(x_i) &= \left(\bigotimes_{j \in \partial I \setminus i} q_{j \rightarrow I}(\emptyset) \right) \otimes \left(\bigoplus_{\underline{x}_I} f_I(\underline{x}_I) \bigotimes_{j \in \partial I \setminus i} \hat{\underline{p}}_{j \rightarrow I}(x_j) \right) \Rightarrow \\ q_{I \rightarrow i}(\emptyset) &= \tilde{\underline{p}}_{I \rightarrow i}(\emptyset) \bigotimes_{j \in \partial I \setminus i} q_{j \rightarrow I}(\emptyset) \end{aligned} \quad (45)$$

Eqs (44) and (45) are simply recursive integration on a tree, where the integral up to node i (i.e., $q_{i \rightarrow I}(\emptyset)$ in eq (44)) is reduced to the integral over its sub-trees. By unrolling this recursion we see that $q_{i \rightarrow I}(\emptyset)$ is simply the product of all $\tilde{\underline{p}}_{I \rightarrow i}(\emptyset)$ and $\tilde{\underline{p}}_{I \rightarrow i}(\emptyset)$ in its sub-tree, where the messages are towards the root. Eq (39) tells us that the global integral is not different. Therefore, eqs (44) and (45) we can completely expand the recursion for the global integral. For this, let $\uparrow i$ restrict the ∂i to the factor that is

higher than variable i in the tree (*i.e.*, closer to the root r). Similarly let $\uparrow I$ be the variable that is closer to the root than I . We can write the global integral as

$$q(\emptyset) = \bigotimes_{i, I=\uparrow i} \tilde{p}_{i \rightarrow I}(\emptyset) \bigotimes_{I, i=\uparrow I} \tilde{p}_{I \rightarrow i}(\emptyset) \quad (46)$$

Proposition 5.3 shows that these local integrals can be written in terms of local integrals of interest – *i.e.*,

$$\tilde{p}_{I \rightarrow i}(\emptyset) = \frac{\tilde{p}_I(\emptyset)}{\tilde{p}_{i \leftrightarrow I}(\emptyset)} \quad \text{and} \quad \tilde{p}_{i \rightarrow I}(\emptyset) = \frac{\tilde{p}_i(\emptyset)}{\tilde{p}_{i \leftrightarrow I}(\emptyset)}$$

Substituting from the equations above into eq (46) we get the equations of theorem 5.2. $\square$

5.2. The new factor-graph and semiring

The decomposition of integral in theorem 5.2 means $Q(\hat{p}_{\rightarrow \cdot})(\emptyset)$ has a factored form. Therefore, a factor-graph with $\hat{p}_{\rightarrow \cdot}$ as the set of variables and three different types of factors corresponding to different terms in the decomposition – *i.e.*, $\tilde{P}_I(\hat{p}_{\partial I \rightarrow I})(\emptyset)$, $\tilde{P}_i(\hat{p}_{\partial i \setminus I \rightarrow i})(\emptyset)$ and $\tilde{P}_{i \leftrightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i})(\emptyset)^{-1}$ – can represent $Q(\hat{p}_{\rightarrow \cdot})(\emptyset)$.

Figure 3 shows a simple factor-graph and the corresponding SP factor-graph. The new factor-graph has one variable per each message in the original factor-graph and three types of factors as discussed above. Survey propagation is simply belief propagation applied to the this new factor-graph using the new semiring. As before, BP messages are exchanged between variables and factors. But here, we can simplify BP messages by substitution and only keep two types of factor-to-factor messages. We use $S_{i \rightarrow I}$ and $S_{I \rightarrow i}$ to denote these two types of SP messages. These messages are exchanged between two types of factors, namely $\tilde{P}_I(\hat{p}_{\partial I \rightarrow I})(\emptyset)$ and $\tilde{P}_i(\hat{p}_{\partial i \setminus I \rightarrow i})(\emptyset)$. Since the third type of factors $\tilde{P}_{i \leftrightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i})(\emptyset)^{-1}$ are always connected to only two variables, $\hat{p}_{i \rightarrow I}$ and $\hat{p}_{I \rightarrow i}$, we can simplify their role in the SP message update to get

$$S_{i \rightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i}) \propto \bigoplus_{\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i}} \left(\frac{\tilde{P}_i(\hat{p}_{\partial i \setminus I \rightarrow i})(\emptyset)}{\tilde{P}_{i \leftrightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i})(\emptyset)} \bigotimes_{J \in \partial i \setminus I} S_{J \rightarrow i}(\hat{p}_{J \rightarrow I}, \hat{p}_{I \rightarrow J}) \right) \quad (47)$$

$$S_{I \rightarrow i}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i}) \propto \bigoplus_{\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i}} \left(\frac{\tilde{P}_I(\hat{p}_{\partial I \rightarrow I})(\emptyset)}{\tilde{P}_{i \leftrightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i})(\emptyset)} \bigotimes_{j \in \partial I \setminus i} S_{j \rightarrow I}(\hat{p}_{j \rightarrow I}, \hat{p}_{I \rightarrow j}) \right) \quad (48)$$

where in all cases we are assuming the messages $\hat{p}_{\rightarrow \cdot} \in \mathcal{P}$ are consistent with each other – *i.e.*, satisfy BP equations on the original factor-graph. Note that, here again we are using the normalized BP message update and the normalization factor is hidden using $\propto$ sign. This is possible because we assumed $\otimes$ has an inverse. We can further simplify this update using the following proposition.

PROPOSITION 5.3. For $\hat{p}_{\rightarrow \cdot} \in \mathcal{P}$

$$\frac{\tilde{P}_i(\hat{p}_{\partial i \setminus I \rightarrow i})(\emptyset)}{\tilde{P}_{i \leftrightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i})(\emptyset)} = \tilde{P}_{i \rightarrow I}(\hat{p}_{\partial i \setminus I \rightarrow i})(\emptyset) \quad (49)$$

and

$$\frac{\tilde{P}_I(\hat{p}_{\partial I \rightarrow I})(\emptyset)}{\tilde{P}_{i \leftrightarrow I}(\hat{p}_{i \rightarrow I}, \hat{p}_{I \rightarrow i})(\emptyset)} = \tilde{P}_{I \rightarrow i}(\hat{p}_{\partial I \setminus i \rightarrow I})(\emptyset) \quad (50)$$




Fig. 3. Part of a factor-graph (left) and the corresponding SP factor-graph on the right. The variables in SP factor-graph are the messages in the original graph. The SP factor-graph has three type of factors: (I) $\tilde{P}_I(\cdot)(\emptyset)$, (II) $\tilde{P}_i(\cdot)(\emptyset)$ and (III) $\tilde{P}_{i \leftrightarrow I}(\cdot)(\emptyset)^{-1}$. As the arrows suggest, SP message updates are simplified so that only two type of messages are exchanged: $S_{i \rightarrow I}$ and $S_{I \rightarrow i}$ between factors of type (I) and (II).

PROOF. By definition of $\tilde{p}_I(\underline{x}_I)$ and $\hat{p}_{i \rightarrow I}(x_i)$ in eqs (32) and (34)

$$\begin{aligned} \tilde{p}_I(x_i) &= \tilde{p}_{I \rightarrow i}(x_i) \otimes \hat{p}_{i \rightarrow I}(x_i) \Rightarrow \bigoplus_{x_i} \tilde{p}_I(x_i) = \bigoplus_{x_i} \tilde{p}_{I \rightarrow i}(x_i) \otimes \hat{p}_{i \rightarrow I}(x_i) \Rightarrow \\ \tilde{p}_I(\emptyset) &= \tilde{p}_{I \rightarrow i}(\emptyset) \otimes \left(\bigoplus_{x_i} \hat{p}_{i \rightarrow I}(x_i) \otimes \hat{p}_{i \rightarrow I}(x_i) \right) \Rightarrow \tilde{p}_I(\emptyset) = \tilde{p}_{I \rightarrow i}(\emptyset) \otimes \tilde{p}_{i \leftrightarrow I}(\emptyset) \end{aligned}$$

where in the last step we used eq (36).

Similarly for the second statement of the proposition we have

$$\begin{aligned} \tilde{p}_i(x_i) &= \tilde{p}_{i \rightarrow I}(x_i) \otimes \hat{p}_{I \rightarrow i}(x_i) \Rightarrow \bigoplus_{x_i} \tilde{p}_i(x_i) = \bigoplus_{x_i} \tilde{p}_{i \rightarrow I}(x_i) \otimes \hat{p}_{I \rightarrow i}(x_i) \Rightarrow \\ \tilde{p}_i(\emptyset) &= \tilde{p}_{i \rightarrow I}(\emptyset) \otimes \left(\bigoplus_{x_i} \hat{p}_{I \rightarrow i}(x_i) \otimes \hat{p}_{I \rightarrow i}(x_i) \right) \Rightarrow \tilde{p}_i(\emptyset) = \tilde{p}_{i \rightarrow I}(\emptyset) \otimes \tilde{p}_{i \leftrightarrow I}(\emptyset) \end{aligned}$$

□

The term on the l.h.s. in the proposition above appear in eqs (47) and (48) and the terms on the r.h.s. are local message integrals given by eqs (32) and (33). We can enforce $\hat{p}_{\cdot \rightarrow \cdot} \in \mathcal{P}$, by enforcing BP updates $\hat{p}_{i \rightarrow I} = P_{i \rightarrow I}(\hat{p}_{\partial i \setminus I \rightarrow i})$ and $\hat{p}_{I \rightarrow i} = P_{I \rightarrow i}(\hat{p}_{\partial I \setminus i \rightarrow I})$ “locally”, during the message updates in the new factor-graph. Combining this constraint with the simplification offered by proposition 5.3 gives us the SP message updates

$$S_{i \rightarrow I}(\hat{p}_{i \rightarrow I}) \propto \bigoplus_{\hat{p}_{\partial i \setminus I \rightarrow i}} \left(1(\hat{p}_{i \rightarrow I} = P_{i \rightarrow I}(\hat{p}_{\partial i \setminus I \rightarrow i})) \otimes \tilde{P}_{i \rightarrow I}(\hat{p}_{\partial i \setminus I \rightarrow i})(\emptyset) \otimes_{j \in \partial i \setminus I} S_{j \rightarrow i}(\hat{p}_{j \rightarrow i}) \right) \quad (51)$$

$$S_{I \rightarrow i}(\hat{p}_{I \rightarrow i}) \propto \bigoplus_{\hat{p}_{\partial I \setminus i \rightarrow I}} \left(1(\hat{p}_{I \rightarrow i} = P_{I \rightarrow i}(\hat{p}_{\partial I \setminus i \rightarrow I})) \otimes \tilde{P}_{I \rightarrow i}(\hat{p}_{\partial I \setminus i \rightarrow I})(\emptyset) \otimes_{j \in \partial I \setminus i} S_{j \rightarrow I}(\hat{p}_{j \rightarrow I}) \right) \quad (52)$$

where $1(\cdot)$ is the identity function on the SP semiring, where $1(\text{TRUE}) = \overset{\otimes}{1}$ and $1(\text{FALSE}) = \overset{\oplus}{1}$.

Here each SP message is a functional over all possible BP messages between the same variable and factor. However, in updating the SP messages, the identity functions

ensure that only the messages that locally satisfy BP equations are taken into account. Another difference from the updates of eqs (47) and (48) is that SP messages have a single argument. This is because the new local integrals either depend on $\hat{p}_{i \rightarrow I}$ or $\hat{p}_{I \rightarrow i}$, and not both.

Example 5.4. In variational approach, survey propagation comes in two variations: entropic SP(ξ) and energetic SP(y) [Mézard and Montanari 2009]. For the readers familiar with variational derivation of SP, here we express the relation to the algebraic approach. According to the variational view, the partition function of the *entropic SP* is $\sum_{\hat{p}_{\rightarrow}} e^{\xi \log(Q(\hat{p}_{\rightarrow})(\emptyset))}$, where $Q(\hat{p}_{\rightarrow})(\emptyset)$ is the partition function for the sum-product semiring. The entropic SP has an inverse temperature parameter, *a.k.a. Parisi parameter*, $\xi \in \mathbb{R}$. It is easy to see that $\xi = 1$ corresponds to $\hat{\oplus} = +$, $\hat{\oplus} = +$ and $\hat{\otimes} = \times$ in our algebraic approach. The limits of $\xi \rightarrow \infty$ corresponds to $\hat{\oplus} = \max$. On the other hand, the limit of $\xi \rightarrow 0$ amounts to ignoring $Q(\hat{p}_{\rightarrow})(\emptyset)$ and corresponds to the counting SP; see section 5.4.

The *energetic SP*(y) is different only in the sense that $Q(\hat{p}_{\rightarrow})(\emptyset)$ in $\sum_{\hat{p}_{\rightarrow}} e^{-y \log(Q(\hat{p}_{\rightarrow})(\emptyset))}$ is the ground state energy. This corresponds to $\hat{\oplus} = +$, $\hat{\oplus} = \max$ and $\hat{\otimes} = \sum$, and the limits of the inverse temperature parameter $y \rightarrow \infty$ is equivalent to $\hat{\oplus} = \min$, $\hat{\oplus} = \min$ and $\hat{\otimes} = \sum$. By taking an algebraic view we can choose between both operations and domains. For instance, an implication of algebraic view is that all the variations of SP can be applied to the domain of complex numbers $\mathcal{Y}^* = \mathbb{C}$.

5.3. The new integral and marginals

Once again we can use theorem 5.2, this time to approximate the *SP integral* $Q(\emptyset)(\emptyset) = \hat{\bigoplus}_{\hat{p}_{\rightarrow}} Q(\hat{p}_{\rightarrow})(\emptyset)$ using local integral of SP messages.

The *SP marginal* over each BP message $\hat{p}_{i \rightarrow I}$ or $\hat{p}_{I \rightarrow i}$ is the same as the corresponding SP message – i.e., $S(\hat{p}_{i \rightarrow I}) = S_{i \rightarrow I}(\hat{p}_{i \rightarrow I})$. To see this in the factor-graph of figure 3, note that each message variable is connected to two factors, and both of these factors are already contained in calculating one SP messages.

Moreover, from the SP marginals over messages we can recover the SP marginals over BP marginals which we denote by $S(\hat{p})(x_i)$. For this, we simply need to enumerate all combinations of BP messages that produce a particular marginal (weighting them by their local integral $\tilde{P}_i(\hat{p}_{\partial i \rightarrow i})(\emptyset)$)

$$S(\hat{p})(x_i) \propto \hat{\bigoplus}_{\hat{p}_{\partial i \rightarrow i}} 1(\hat{p}(x_i) = P(\hat{p}_{\partial i \rightarrow i})(x_i)) \otimes \tilde{P}_i(\hat{p}_{\partial i \rightarrow i})(\emptyset) \bigotimes_{I \in \partial i} S_{I \rightarrow i}(\hat{p}_{I \rightarrow i}) \quad (53)$$

5.4. Counting survey propagation

Previously we required the $\otimes$ operator to have an inverse, so that we can decompose the BP integral $q(\emptyset)$ into local integrals. Moreover, for a consistent decomposition of the BP integral, SP and BP semiring previously shared the $\otimes$ operation.⁸

Here, we lift these requirements by discarding the BP integrals altogether. This means SP semiring could be completely distinct from BP semiring and $(\mathcal{Y}^*, \otimes)$ does not have to be an Abelian group. This setting is particularly interesting when the SP

⁸This is because, if the expansion operation $\hat{\otimes}$ was different from the expansion operation of BP, $\otimes$, the expanded form $Q(\hat{p}_{\rightarrow})$ in the SP factor-graph would not evaluate the integral $q(\emptyset)$ in the BP factor-graph, even in factor-graphs without any loops.

semiring is sum-product over real domain

$$S_{i \rightarrow I}(\hat{p}_{i \rightarrow I}) \propto \sum_{\hat{p}_{\partial i \setminus I \rightarrow i}} 1(\hat{p}_{i \rightarrow I} = P_{i \rightarrow I}(\hat{p}_{\partial i \setminus I \rightarrow i})) \prod_{J \in \partial i \setminus I} S_{J \rightarrow i}(\hat{p}_{J \rightarrow i}) \quad (54)$$

$$S_{I \rightarrow i}(\hat{p}_{I \rightarrow i}) \propto \sum_{\hat{p}_{\partial I \setminus i \rightarrow I}} 1(\hat{p}_{I \rightarrow i} = P_{I \rightarrow i}(\hat{p}_{\partial I \setminus i \rightarrow I})) \prod_{j \in \partial I \setminus i} S_{j \rightarrow I}(\hat{p}_{j \rightarrow I}) \quad (55)$$

Here, the resulting SP integral $Q(\hat{p}_{\rightarrow}) = \sum_{\hat{p}_{\rightarrow}} 1(\hat{p}_{\rightarrow} \in \mathcal{P})$ simply “counts” the number of BP fixed points and SP marginals over BP marginals (given by eq (53)) which approximates the frequency of a particular marginal. The original survey propagation equations in [Braunstein et al. 2002a], which are very successful in solving satisfiability, correspond to counting SP applied to the or-and semiring.

Example 5.5. Interestingly, in all min-max problems with discrete domains $\mathcal{X}$, min-max BP messages can only take the values that are in the range of factors – i.e., $\mathcal{Y}^* = \mathcal{Y}$. This is because any ordered set is closed under min and max operations. Here, each counting SP message $S_{i \rightarrow I}(\hat{p}_{i \rightarrow I}) : \mathcal{Y}^{|\mathcal{X}_i|} \rightarrow \mathbb{R}$ is a discrete distribution over all possible min-max BP messages. This means counting survey propagation where the BP semiring is min-max is computationally “tractable”. In contrast, (counting) SP, when applied to sum-product BP over real domains, is not tractable. This is because, in this case, each SP message is a distribution over an uncountable set: $S_{i \rightarrow I}(\hat{p}_{i \rightarrow I}) : \mathbb{R}^{|\mathcal{X}_i|} \rightarrow \mathbb{R}$.

In practice, (counting) SP is only interesting if it remains tractable. The most well-known case corresponds to counting SP when applied to the or-and semiring. In this case the factors are constraints and the domain of SP messages is $\{\text{TRUE}, \text{FALSE}\}^{|\mathcal{X}_i|}$. Our algebraic perspective extends this set of tractable instances. For example, it shows that counting SP can be used to count the number of fixed points of BP when applied to xor-and or min-max semiring.

Conclusion

This paper builds on previous work to address three basic questions about inference in graphical models: **(I)** “*what is an inference problem in a graphical model?*” We use the combination of commutative semigroups and a factor-graph to answer this question in a broad sense, generalizing a variety of previous models. **(II)** “*How difficult is inference?*” By confining inference to four operations of min, max, sum and product that easily lend themselves to models of computation, we build an inference hierarchy that is complete for PSPACE and organizes inference problems into complexity classes with increasing levels of difficulty. Only a few of these problems had previously been studied and only a handful of them now have a variational interpretation. Moreover, we prove that inference for “any” commutative semiring is NP-hard under randomized reduction, which generalizes previous results for particular semirings. **(III)** “*When does distributive law help?*” After reviewing the algebraic form of belief propagation, and the conditions that allow its normalized form, we show that application of distributive law in performing exact inference is limited to inference problems with one marginalization operation. **(IV)** Finally we extend the algebraic treatment of message passing techniques to survey propagation. This perspective generalizes SP to new settings and we hope that it also makes this powerful message passing procedure – that has its origins in statistical physics – more accessible to the machine learning community.

REFERENCES

- S Aji and R McEliece. 2001. The Generalized distributive law and free energy minimization. In *Allerton Conf.*

- Srinivas M Aji, Gavin B Horn, and Robert J McEliece. 1998. On the Convergence of iterative decoding on graphs with a single cycle. In *Proc 1998 IEEE Int Symp Information Theory*.
- S M Aji and R J McEliece. 2000. The generalized distributive law. *Information Theory, IEEE Transactions on* 46, 2 (2000), 325–343.
- Sanjeev Arora and Boaz Barak. 2009. *Computational complexity: a modern approach*. Cambridge University Press.
- Fahiem Bacchus. 1991. *Representing and reasoning with probabilistic knowledge: a logical approach to probabilities*. MIT press.
- Mohsen Bayati, Devavrat Shah, and Mayank Sharma. 2005. Maximum weight matching via max-product belief propagation. In *ISIT*. IEEE, 1763–1767.
- Stefano Bistarelli, Ugo Montanari, Francesca Rossi, Thomas Schiex, Gérard Verfaillie, and Hélène Fargier. 1999. Semiring-based CSPs and valued CSPs: Frameworks, properties, and comparison. *Constraints* 4, 3 (1999), 199–240.
- Lucas Bordeaux and Eric Monfroy. 2002. Beyond NP: Arc-consistency for quantified constraints. In *Principles and Practice of Constraint Programming-CP 2002*. Springer, 371–386.
- A Braunstein, M Mezard, M Weigt, and R Zecchina. 2002b. Constraint Satisfaction by Survey Propagation. *Physics* 2 (2002), 8.
- A Braunstein, M Mezard, and R Zecchina. 2002a. Survey propagation: an algorithm for satisfiability. *Random Structures and Algorithms* 27, 2 (2002), 19.
- A. Braunstein and R. Zecchina. 2003. Survey Propagation as local equilibrium equations. *arXiv:cond-mat/0312483* (Dec. 2003). DOI: <http://dx.doi.org/10.1088/1742-5468/2004/06/P06007> J. Stat. Mech., P06007 (2004).
- Peter Clifford. 1990. Markov random fields in statistics. *Disorder in physical systems* (1990), 19–32.
- G Cooper. 1990. The computational complexity of probabilistic inference using Bayesian belief networks. *Artificial Intelligence* 42 (1990).
- Brian A Davey and Hilary A Priestley. 2002. *Introduction to lattices and order*. Cambridge university press.
- Rina Dechter and David Larkin. 2001. Hybrid processing of beliefs and constraints. In *Proceedings of the Seventeenth conference on Uncertainty in artificial intelligence*. Morgan Kaufmann Publishers Inc., 112–119.
- Beno Eckmann and Peter J Hilton. 1962. Group-like structures in general categories I multiplications and comultiplications. *Math. Ann.* 145, 3 (1962), 227–255.
- G David Forney Jr. 2001. Codes on graphs: normal realizations. *Information Theory, IEEE Transactions on* 47, 2 (2001), 520–548.
- Y Fu and P W Anderson. 1986. Application of statistical mechanics to NP-complete problems in combinatorial optimisation. *Journal of Physics A: Mathematical and General* 19, 9 (June 1986), 1605–1620. DOI: <http://dx.doi.org/10.1088/0305-4470/19/9/033>
- Robert G Gallager. 1962. Low-density parity-check codes. *Information Theory, IRE Transactions on* 8, 1 (1962), 21–28.
- T Heskes. 2003. Stable Fixed Points of Loopy Belief Propagation Are Local Minima of the Bethe Free Energy. In *NIPS*.
- Ronald A Howard and James E Matheson. 2005. Influence diagrams. *Decision Analysis* 2, 3 (2005), 127–143.
- Bert Huang and Tony Jebara. 2009. Approximating the permanent with belief propagation. *arXiv preprint arXiv:0908.1769* (2009).
- Morteza Ibrahimi, Adel Javanmard, Yashodhan Kanoria, and Andrea Montanari. 2011. Robust max-product belief propagation. In *Signals, Systems and Computers (ASILOMAR), 2011 Conference Record of the Forty Fifth Asilomar Conference on*. IEEE, 43–49.
- Edwin T Jaynes. 2003. *Probability theory: the logic of science*. Cambridge university press.
- Michael Kearns. 2007. Graphical games. *Algorithmic game theory* 3 (2007), 159–180.
- Lukas Kroc, A Sabharwal, and Bart Selman. 2002. Survey Propagation Revisited. *23rd UAI* (2002), 217–226.
- F R Kschischang and B J Frey. 2001. Factor graphs and the sum-product algorithm. *Information Theory, IEEE* (2001).
- Steffen L Lauritzen and Finn Verner Jensen. 1997. Local computation with valuations from a commutative semigroup. *Annals of Mathematics and Artificial Intelligence* 21, 1 (1997), 51–69.
- Michael L Littman, Stephen M Majercik, and Toniann Pitassi. 2001. Stochastic boolean satisfiability. *Journal of Automated Reasoning* 27, 3 (2001), 251–296.
- E N Maneva, E Mossel, and M J Wainwright. 2004. A New Look at Survey Propagation and its Generalizations. *arXiv:cs/0409012* (Sept. 2004).

- M. Mézard and A. Montanari. 2009. *Information, physics, and computation*. Oxford.
- Marc Mézard, Giorgio Parisi, and M. A. Virasoro. 1987. *Spin Glass Theory and Beyond*. Singapore: World Scientific.
- M. Mezard, G. Parisi, and R. Zecchina. 2002. Analytic and Algorithmic Solution of Random Satisfiability Problems. *Science* 297, 5582 (Aug. 2002), 812–815. DOI: <http://dx.doi.org/10.1126/science.1073287>
- D Mitchell, B Selman, and H Levesque. 1992. Hard and easy distributions of SAT problems. In *AAAI*, Vol. 92. 459–465.
- Rémi Monasson, Riccardo Zecchina, Scott Kirkpatrick, Bart Selman, and Lidror Troyansky. 1999. Determining computational complexity from characteristic phase transitions. *Nature* 400, 6740 (1999), 133–137.
- Kevin Murphy, Yair Weiss, and Michael Jordan. 1999. Loopy-belief Propagation for Approximate Inference: An Empirical Study. In 15. 467–475.
- Luis E Ortiz and Michael Kearns. 2002. Nash propagation for loopy graphical games. In *Advances in Neural Information Processing Systems*. 793–800.
- James D Park and Adnan Darwiche. 2004. Complexity Results and Approximation Strategies for MAP Explanations. *J. Artif. Intell. Res.(JAIR)* 21 (2004), 101–133.
- Judea Pearl. 1985. Bayesian networks: a model of self-activated memory for evidential reasoning. (1985).
- J Pearl. 1988. *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*. Representation and Reasoning, Vol. 88. Morgan Kaufmann.
- Charles C Pinter. 2012. *A book of abstract algebra*. Courier Dover Publications.
- Brian Potetz and Tai Sing Lee. 2008. Efficient Belief Propagation for Higher Order Cliques Using Linear Constraint Nodes Cliques Using Linear Constraint Nodes. *Computer Vision and Image Understanding* 112, 1 (2008), 39–54. DOI: <http://dx.doi.org/10.1016/j.cviu.2008.05.007>
- Cédric Pralet, Gérard Verfaillie, and Thomas Schiex. 2007. An Algebraic Graphical Model for Decision with Uncertainties, Feasibilities, and Utilities. *J. Artif. Intell. Res.(JAIR)* 29 (2007), 421–489.
- S Ravanbakhsh, C Srinivasa, B Frey, and R Greiner. 2014. Min-Max Problems on Factor-Graphs. *ICML* (2014).
- Hartley Rogers. 1987. Theory of recursive functions and effective computability. (1987).
- Dan Roth. 1993. On the hardness of approximate reasoning. In *Proceedings of the 13th international joint conference on Artificial intelligence-Volume 1*. Morgan Kaufmann Publishers Inc., 613–618.
- Thomas Schiex, Helene Fargier, and Gerard Verfaillie. 1995. Valued constraint satisfaction problems: hard and easy problems. In *Proceedings of the 14th international joint conference on Artificial intelligence-Volume 1*. Morgan Kaufmann Publishers Inc., 631–637.
- Prakash P Shenoy. 1992. Valuation-based systems: A framework for managing uncertainty in expert systems. In *Fuzzy logic for the management of uncertainty*. John Wiley & Sons, Inc., 83–104.
- Prakash P Shenoy and Glenn Shafer. 1990. Axioms for probability and belief-function propagation. In *Uncertainty in Artificial Intelligence*.
- Solomon Eyal Shimony. 1994. Finding MAPs for belief networks is NP-hard. *Artificial Intelligence* 68, 2 (1994), 399–410.
- Larry J Stockmeyer. 1976. The polynomial-time hierarchy. *Theoretical Computer Science* 3, 1 (1976), 1–22.
- Daniel Tarlow and Richard S Zemel. 2010. HOP-MAP : Efficient Message Passing with High Order Potentials. *Journal of Machine Learning Research Proceedings Track 9* (2010), 812–819. http://www.cs.toronto.edu/~dtarlow/TarlowGivoniZemel_AISTATS2010.pdf
- Seinosuke Toda. 1991. PP is as hard as the polynomial-time hierarchy. *SIAM J. Comput.* 20, 5 (1991), 865–877.
- L G. Valiant. 1979. The complexity of computing the permanent. *Theoretical computer science* 8, 2 (1979), 189–201.
- Leslie G Valiant and Vijay V Vazirani. 1986. NP is as easy as detecting unique solutions. *Theoretical Computer Science* 47 (1986), 85–93.
- John Von Neumann and Oskar Morgenstern. 2007. *Theory of Games and Economic Behavior (60th Anniversary Commemorative Edition)*. Princeton university press.
- Klaus W Wagner. 1986. The complexity of combinatorial problems with succinct input representation. *Acta Informatica* 23, 3 (1986), 325–356.
- Martin J Wainwright and Michael I Jordan. 2007. Graphical Models, Exponential Families, and Variational Inference. *Foundations and Trends in Machine Learning* 1, 12 (2007), 1–305. DOI: <http://dx.doi.org/10.1561/22000000001>
- Yair Weiss. 2001. Correctness of belief propagation in Gaussian graphical models of arbitrary topology. *Neural computation* June (2001).

- Adrian Weller and Tony S Jebara. 2013. On MAP inference by MWSS on Perfect Graphs. *arXiv preprint arXiv:1309.6872* (2013).
- JS Yedidia and WT Freeman. 2001. Generalized belief propagation. *Advances in neural information* (2001).